hep-th/9604105

The Non-Critical $N = 2$ String is an $s\ell(2|1)$ Theory

A. M. Semikhatov

I. E. Tamm Theory Division, P. N. Lebedev Physics Institute, Russian Academy of Sciences

arXiv:hep-th/9604105v3 15 Jan 1997

Abstract

Any non-critical $N=2$ string is shown to have an affine $s\ell(2|1)$ worldsheet symmetry in the *conformal* gauge.

1 Introduction

$N=2$ strings [1] have been studied now for about two decades (see [2] for a review). The $(2,2)$ theory was first quantized a la Polyakov in [3], where the critical dimension was found to be 4. In the BRST approach, this result was obtained and further discussed in [4, 5]. That the target space is *complex two* dimensional was realized in [6], where the geometry for the $N=2$ strings was investigated and shown to be self-dual. The moduli and BRST cohomology of the $N=2$ strings were studied in [7]. Recently, a very important role was proposed for the heterotic $(2,1)$ strings in the M-theory [8].

Non-critical string theories involve, in addition to the corresponding (super)matter theory and ghosts, the (super)Liouville sector whose central charge adds up with the matter central charge to the critical value. The $N=2$ string is described on the worldsheet by $N=2$ matter coupled to two dimensional $N=2$ supergravity [9]. This string theory is singled out for a number of reasons, one of them being that the bosonic ($N=0$) and $N=1$ strings can be embedded into it [10, 11, 12, 13, 14]. On the other hand, the $N=0,1,2$ non-critical string theories in the conformal gauge have hidden $N+2$ twisted supersymmetry algebras on the worldsheet [15, 16, 17]. The realizations of the $N+2$ algebras provided by the non-critical string theories can be obtained via hamiltonian reduction from the affine Lie superalgebras [16, 18]. This gives rise to the proposal [17, 19, 20] to classify string theories in terms of Kač–Moody (super)algebras, the idea not unrelated to the embeddings of different string theories into each other [11].

The various relations between non-critical string theories and WZW models have been studied from different points of view for a long time, and the $N+2$ *supersymmetry* algebras are in fact a step towards the WZW theories. The simplest case corresponding to the bosonic string is the affine $s\ell(2)$ algebra; the $N=2$ superconformal algebra, which is realized in the bosonic string, does indeed have many features in common with $s\ell(2)$ (thus, the two algebras 'share' the parafermions and have intimately related singular vectors and in fact similar representation theories [21, 22]). Moreover, the realization of the $N=2$ superconformal algebra in the bosonic string carries over to an $s\ell(2)$ representation in terms of an arbitrary matter theory tensored with several free-field theories [23]. This pattern extends to the $N=1$ superstring and the $osp(1|2)$ Kač–Moody algebra [24]. Thus, in the two lowest cases ($N=0$ and $N=1$), by tensoring the respective non-critical string with an additional free scalar, one can reconstruct the ambient space of the hamiltonian reduction: the corresponding Kač–Moody currents and the 'Drinfeld–Sokolov' ghosts (those that are needed in order to build the BRST complex for the hamiltonian reduction).

In this paper we show that for the non-chiral $N=2$ (i.e., $(2,2)$) string this reconstruction of, in that case, the $s\ell(2|1)$ Kač–Moody superalgebra can always be achieved directly in the space of fields of the non-critical $N=2$ string in the conformal gauge, irrespective of the realization of its matter sector. Therefore, any non-critical $N=2$ string bears a representation of the $s\ell(2|1)$ current algebra. In fact, at the level of comparing the energy-momentum tensors, the non-critical $N=2$ string is the same as the $s\ell(2|1)$ current algebra together with two fermionic and two bosonic first-order systems, the 'Drinfeld–Sokolov' ghosts.

Thus, while the hamiltonian reduction of $s\ell(2|1)$ gives rise to the $N = 2$ superconformal algebra [25, 16, 18], dressing that 'matter' theory with the $N=2$ gravity multiplet allows us to 'invert' the reduction, i.e. to reconstruct both the $s\ell(2|1)$ currents and the Drinfeld–Sokolov ghosts.

In section 2, we give the main construction of this paper, the $s\ell(2|1)$ currents realized in any non-critical $N = 2$ string. Restricting this construction to the $osp(1|2)$ and $s\ell(2)$ subalgebras will result in a new pattern of mappings between $N = 2, 1, 0$ non-critical strings, which we consider in section 3. An adaptation of the new realization of $s\ell(2|1)$ for the topological string is given in section 4. In section 5 we consider the representation spaces and construct the $s\ell(2|1)$ highest-weight states in the tensor product of certain $N=2$ matter Verma modules with free field modules, and outline the corresponding KPZ-like formulation (which appears to have become tautological by virtue of our construction of $s\ell(2|1)$). In section 6 we consider a 'non-stringy', more economical, reformulation of the new realization of $s\ell(2|1)$ in terms of $N = 2$ matter dressed with free fields, the latter no longer being those of the $N = 2$ string; relation of the new realization to the Wakimoto representations of $s\ell(2|1)$ is also considered here. Section 7 contains several concluding remarks, and the Appendix summarizes our conventions.

2 $s\ell(2|1)$ in the $N=2$ supergravity

In this section we will present our main construction, a representation of the affine $s\ell(2|1)$ algebra (A.1) in terms of the fields of the $N=2$ noncritical string. The $N = 2$ matter is described by the energy-momentum tensor $T_{\rm m}(z)$, the $U(1)$ current $H_{\rm m}(z)$ and two supersymmetry currents $G_{\rm m}(z)$ and $\overline{G}_{\rm m}(z)$. The operator products are written out in the Appendix. The $N=2$ matter central charge is

$$c_{\rm m} = 3 - 6q^2 = -3 - 6k\,, \tag{2.1}$$

where the matter background charge q^2 satisfies[1] $q^2 = k + 1$.

In the non-critical $N=2$ string one also has the reparametrization ghosts bc and the $U(1)$ ghosts $\eta\xi$:

$$b(z)\,c(w) = \frac{1}{z-w}\,, \qquad \eta(z)\,\xi(w) = \frac{1}{z-w}\,, \tag{2.2}$$

and the bosonic ghosts

$$\widetilde{\beta}(z)\,\widetilde{\gamma}(w) = \frac{-1}{z-w}\,, \qquad \beta(z)\,\gamma(w) = \frac{-1}{z-w}\,, \tag{2.3}$$

and, finally, a super-Liouville sector represented in components by a free complex scalar and a complex fermion:

$$\partial\phi(z)\,\partial\overline{\phi}(w) = \frac{-1}{(z-w)^2}\,, \qquad \psi(z)\,\overline{\psi}(w) = \frac{1}{z-w}\,. \tag{2.4}$$

The energy-momentum tensor reads

$$\begin{aligned}\mathsf{T} \;=\; & T_{\rm m} - \partial bc - 2b\partial c - \eta\partial\xi - \tfrac{1}{2}\partial\beta\gamma - \tfrac{3}{2}\beta\partial\gamma - \tfrac{1}{2}\partial\widetilde{\beta}\widetilde{\gamma} - \tfrac{3}{2}\widetilde{\beta}\partial\widetilde{\gamma}\\ & - \partial\phi\partial\overline{\phi} + \tfrac{1}{2}\,\partial\partial\overline{\phi} + \tfrac{1}{2}q^2\,\partial\partial\phi + \tfrac{1}{2}\partial\psi\overline{\psi} - \tfrac{1}{2}\psi\partial\overline{\psi}\end{aligned} \tag{2.5}$$

We do not make any assumptions on the nature of the $N=2$ matter represented by the energy-momentum tensor $T_{\rm m}$, the $U(1)$ current $H_{\rm m}$ and the supercurrents $G_{\rm m}$ and $\overline{G}_{\rm m}$. This can be any such theory, hence our construction is similar to those of refs. [15, 16, 23, 17] in that the matter sector is only required to satisfy the appropriate (super)conformal algebra. Another remark is purely technical: the OPEs (2.4) are invariant under rescalings

---

[1] These formulae simply introduce a new parameter, k; later this will become the level of the $s\ell(2|1)$ algebra we are going to construct. The matter central charge will thus be unambiguously related to the level.

$$\phi \mapsto a\phi\,, \qquad \overline{\phi} \mapsto a^{-1}\overline{\phi}\,, \tag{2.6}$$

therefore only the product $\overline{Q}Q$ of the Liouville background charges introduced into the energy-momentum tensor as $+\frac{1}{2}\,\overline{Q}\partial\partial\overline{\phi} + \frac{1}{2}Q\partial\partial\phi$ has an invariant meaning. The central charge in the super-Liouville sector is $3 + 6\overline{Q}Q = 3 + 6q^2$.

Now we are going to use the fields $T_{\rm m}$, $H_{\rm m}$, $\overline{G}_{\rm m}$, $G_{\rm m}$, b, c, η, ξ, $\widetilde{\beta}$, $\widetilde{\gamma}$, β, γ, $\partial\phi$, $\partial\overline{\phi}$, ψ, $\overline{\psi}$ to construct the $s\ell(2|1)$ currents E_1, E_2, E_{12}, H^-, H^+, F_1, F_2, F_{12} that would satisfy the OPEs (A.1).

The upper-triangular currents are an easy part:

$$\begin{aligned} E_1 &= \psi\, e^{\phi}\,, \\ E_2 &= \overline{\psi}\, e^{\phi}\,, \\ E_{12} &= e^{2\phi}\,. \end{aligned} \tag{2.7}$$

The choice of ψ vs. $\overline{\psi}$ is merely a convention; also, the exponents in (2.7) may be changed by a common factor due to (2.6).

Now we expect the matter ingredients $T_{\rm m}$, $G_{\rm m}$, $\overline{G}_{\rm m}$, and $H_{\rm m}$ to enter the lower-diagonal currents, i.e., the matter fields should go to where they are to be found after the hamiltonian reduction. We thus expect $T_{\rm m}$ to go into F_{12}. For $\overline{G}_{\rm m}$ and $G_{\rm m}$ we also have natural candidates: F_1 and F_2 respectively or vice versa (the choice is conventional in view of the automorphism of the algebra (A.3) that includes the mapping $\overline{G}_{\rm m} \leftrightarrow G_{\rm m}$). Further, we have H^+ to incorporate the matter $H_{\rm m}$ current. Then, H^- is a linear combination of the remaining currents that one has in the $N{=}2$ non-critical string:

$$\begin{aligned} H^- &= -\tfrac{1}{2}\partial\overline{\phi} + \tfrac{1}{2}(k-3)\partial\phi + b\,c + \tfrac{1}{2}\beta\,\gamma + \tfrac{1}{2}\widetilde{\beta}\,\widetilde{\gamma} + \eta\,\xi \\ H^+ &= -\tfrac{1}{2}H_{\rm m} + \tfrac{1}{2}\psi\,\overline{\psi} \end{aligned} \tag{2.8}$$

These currents are mutually orthogonal, normalized properly and satisfy the required OPEs with the upper triangular currents (2.7) [2].

The precise form of the rest of the algebra is less obvious, considering an enormous number of *potentially* allowed terms. However, these actually reduce to a reasonable amount; thus, F_1 is essentially proportional to $\overline{\psi}$, plus a piece that contains one of the matter supercurrents, as expected:

$$F_1 = \Big(G_{\rm m} - bc\overline{\psi} - \tfrac{1}{2}\beta\gamma\,\overline{\psi} - \tfrac{1}{2}\widetilde{\beta}\widetilde{\gamma}\,\overline{\psi} + \tfrac{1}{2}\partial\overline{\phi}\,\overline{\psi} + \tfrac{1}{2}(3+k)\partial\phi\,\overline{\psi} - \eta\xi\overline{\psi} - \tfrac{1}{2}H_{\rm m}\overline{\psi} - (k+\tfrac{1}{2})\partial\overline{\psi}\Big)e^{-\phi} \tag{2.9}$$

Note that there is a symmetry

$$\overline{G}_{\rm m} \to a\overline{G}_{\rm m}\,, \quad G_{\rm m} \to a^{-1}G_{\rm m} \tag{2.10}$$

of the $N{=}2$ algebra, which allows us to introduce an arbitrary non-zero parameter in front of $G_{\rm m}$ in (2.9) (and then, accordingly, in front of $\overline{G}_{\rm m}$ in the subsequent formulae).

In (2.9), the notation is somewhat abused in the following manner. *Throughout the paper, the nested normal orderings are assumed from right to left, as* $:\!A\,:\!BC\!:\,:$. However, in order to make the formulae shorter, we write the vertex operators $\exp(a\phi)$ as common factors. Thus eq. (2.9) and similar formulae below should be understood by multiplying every term in the parentheses with $e^{-\phi}$ and introducing the normal ordering as explained.

---

[2] We have chosen the 'real' basis for the two $\beta\gamma$ systems rather than the 'complex' basis in which $\beta^{\pm}(z)\,\gamma^{\mp}(w) = -1/(z{-}w)$.

Next, not surprisingly, F_2 is essentially proportional to ψ, plus a term with the other matter supercurrent:

$$F_2 = \Big((k+1)\,\overline{G}_{\rm m} + bc\,\psi + \tfrac{1}{2}\beta\gamma\,\psi + \tfrac{1}{2}\widetilde{\beta}\widetilde{\gamma}\,\psi - \tfrac{1}{2}\partial\overline{\phi}\,\psi - \tfrac{1}{2}(3+k)\partial\phi\,\psi + \eta\xi\,\psi - \tfrac{1}{2}H_{\rm m}\,\psi + (k+\tfrac{1}{2})\partial\psi\Big)e^{-\phi} \quad (2.11)$$

Now F_1 and F_2 give rise to F_{12}:

$$\begin{aligned} F_{12} \;=\; & \Big((k+1)T_{\rm m} - (k+1)\overline{G}_{\rm m}\overline{\psi} + \tfrac{1}{4}H_{\rm m}H_{\rm m} + \tfrac{1}{2}H_{\rm m}\psi\overline{\psi} - bc\beta\gamma - bc\widetilde{\beta}\widetilde{\gamma} + bc\partial\overline{\phi} + (3+k)bc\partial\phi \\ & - 2bc\eta\xi - kb\partial c - \tfrac{1}{4}\beta\beta\gamma\gamma + \tfrac{1}{2}\beta\gamma\partial\overline{\phi} + (\tfrac{3}{2}+\tfrac{k}{2})\beta\gamma\partial\phi - (\tfrac{1}{4}+\tfrac{k}{2})\beta\partial\gamma - \tfrac{1}{4}\widetilde{\beta}\widetilde{\beta}\widetilde{\gamma}\widetilde{\gamma} - \tfrac{1}{2}\widetilde{\beta}\widetilde{\gamma}\beta\gamma \\ & + \tfrac{1}{2}\widetilde{\beta}\widetilde{\gamma}\partial\overline{\phi} + (\tfrac{3}{2}+\tfrac{k}{2})\widetilde{\beta}\widetilde{\gamma}\partial\phi - (\tfrac{1}{4}+\tfrac{k}{2})\widetilde{\beta}\partial\widetilde{\gamma} - \tfrac{1}{4}\partial\overline{\phi}\partial\overline{\phi} - (\tfrac{3}{2}+\tfrac{k}{2})\partial\phi\partial\overline{\phi} - \tfrac{1}{4}(3+k)^2\partial\phi\partial\phi - \eta\xi\beta\gamma \\ & - \eta\xi\widetilde{\beta}\widetilde{\gamma} + \eta\xi\partial\overline{\phi} + (3+k)\eta\xi\partial\phi - k\eta\partial\xi + G_{\rm m}\psi + (\tfrac{1}{4}+\tfrac{k}{2})\psi\partial\overline{\psi} - (2+k)\partial bc - (\tfrac{3}{4}+\tfrac{k}{2})\partial\beta\gamma \\ & - (\tfrac{3}{4}+\tfrac{k}{2})\partial\widetilde{\beta}\widetilde{\gamma} + \tfrac{1}{2}(k+1)\partial\partial\overline{\phi} + \tfrac{1}{2}(3+4k+k^2)\partial\partial\phi - (2+k)\partial\eta\xi - (\tfrac{1}{4}+\tfrac{k}{2})\partial\psi\overline{\psi}\Big)e^{-2\phi} \end{aligned} \quad (2.12)$$

All the $s\ell(2|1)$ OPEs (A.1) are now satisfied (and so are those vanishing ones which are not shown in (A.1) explicitly).

Since the $N=2$ matter is to be recovered as the result of the hamiltonian reduction of $s\ell(2|1)$ [25], the above construction can be viewed as an 'inversion' of the hamiltonian reduction. It is thus achieved by dressing the $N=2$ matter into the $N=2$ string. A natural question is, how much bigger the $N=2$ non-critical string is than the $s\ell(2|1)$ algebra just constructed, that is, what is the 'codimension' of $s\ell(2|1)$ in the $N=2$ string? It turns out that the directions in the $N=2$ string field space orthogonal to the $s\ell(2|1)$ algebra are naturally organized into precisely those ghost fields that have to be introduced when performing the hamiltonian reduction of $s\ell(2|1)$ (including the auxiliary fermionic system). These are two fermionic bc systems and two bosonic $\beta\gamma$ ones; we will call them the *Drinfeld–Sokolov ghosts*, in order to distinguish them from the $N=2$ string ghosts.

The Drinfeld–Sokolov ghosts can be constructed along with the $s\ell(2|1)$ currents out of the same ingredients, the $N=2$ non-critical string, as

$$\begin{aligned} \widetilde{\mathsf{B}} &= \widetilde{\beta}\,e^{-\phi}\,, & \widetilde{\Gamma} &= \widetilde{\gamma}\,e^{\phi}\,, & &\Rightarrow\; \widetilde{\mathsf{B}}(z)\widetilde{\Gamma}(w) = \frac{-1}{z-w} \\ \mathsf{B} &= \beta\,e^{-\phi}\,, & \Gamma &= \gamma\,e^{\phi}\,, & &\Rightarrow\; \mathsf{B}(z)\Gamma(w) = \frac{-1}{z-w} \\ B &= b\,e^{-2\phi}\,, & C &= c\,e^{2\phi}\,, & &\Rightarrow\; B(z)C(w) = \frac{1}{z-w} \\ \mathcal{C} &= \xi\,e^{2\phi}\,, & \mathcal{B} &= \eta\,e^{-2\phi}\,, & &\Rightarrow\; \mathcal{C}(z)\mathcal{B}(w) = \frac{1}{z-w} \end{aligned} \quad (2.13)$$

Then we have the following fundamental identity, which can be interpreted as a 'completeness relation':

$$T_{\rm Sug} + T_{\rm DS} = \mathsf{T}\,, \quad (2.14)$$

where T is the $N=2$ string energy-momentum tensor (2.5), $T_{\rm Sug}$ is the $s\ell(2|1)$ Sugawara energy-momentum tensor (A.2), and the Drinfeld–Sokolov ghost energy-momentum tensor is

$$T_{\rm DS} = -\widetilde{\mathsf{B}}\,\partial\widetilde{\Gamma} - \mathsf{B}\,\partial\Gamma - B\,\partial C - \mathcal{C}\,\partial\mathcal{B}\,. \quad (2.15)$$

The identity (2.14) tells us that the non-critical $N=2$ string is the same, at the level of energy-momentum tensors, as the $s\ell(2|1)$ superalgebra plus the Drinfeld–Sokolov ghosts (we do not consider in this paper the subtle questions of the definitions of the respective spaces of *physical* states and relations between them). Let us note that $\lim_{k\to-1} T_{\rm Sug} = T_{\rm m} + \ldots$ is finite in the realization (2.7)–(2.12), and in fact the identity (2.14) remains valid as $k\to-1$.

In the rest of the paper we consider several immediate consequences of the construction (2.7)–(2.13).

A more compact expression for the $s\ell(2|1)$ currents can be obtained as follows. We introduce a scalar

$$\partial F = -\tfrac{1}{2}\partial\overline{\phi} - \tfrac{1}{2}(3+k)\partial\phi + b\,c + \tfrac{1}{2}\beta\,\gamma + \tfrac{1}{2}\widetilde{\beta}\,\widetilde{\gamma} + \eta\,\xi \tag{2.16}$$

Then the $s\ell(2|1)$ currents rewrite as

$$\begin{aligned}
H^- &= \partial F + k\partial\phi\,,\\
F_1 &= \Bigl(-\overline{\psi}\,\partial F - \tfrac{1}{2}H_{\rm m}\,\overline{\psi} + G_{\rm m} - (k+\tfrac{1}{2})\partial\overline{\psi}\Bigr)e^{-\phi}\,,\\
F_2 &= \Bigl(\psi\,\partial F - \tfrac{1}{2}H_{\rm m}\,\psi + (k+1)\overline{G}_{\rm m} + (k+\tfrac{1}{2})\partial\psi\Bigr)e^{-\phi}\,,\\
F_{12} &= \Bigl(-\partial F\,\partial F - (k+1)\partial\partial F + G_{\rm m}\,\psi + \tfrac{1}{4}H_{\rm m}\,H_{\rm m} + \tfrac{1}{2}H_{\rm m}\,\psi\,\overline{\psi} + (\tfrac{1}{4}+\tfrac{k}{2})\psi\,\partial\overline{\psi}\\
&\quad - (k+1)\overline{G}_{\rm m}\,\overline{\psi} + (k+1)T_{\rm m} + (-\tfrac{1}{4}-\tfrac{k}{2})\partial\psi\,\overline{\psi}\Bigr)e^{-2\phi}
\end{aligned} \tag{2.17}$$

Let us add one more word of caution: as we have remarked, the exponential factors in the above formulae are always assumed to multiply every monomial in the parentheses and the monomials are normal ordered as, e.g., $:\!\partial F\,:\!\partial F\,e^{-2\phi}\!:\,:$.

It is now obvious, in particular, that the bosonic ghosts β, γ, $\widetilde{\beta}$, and $\widetilde{\gamma}$ enter the $s\ell(2|1)$ currents only through their respective ghost currents $:\!\beta\,\gamma\!:$ and $:\!\widetilde{\beta}\,\widetilde{\gamma}\!:$ (clearly, $:\!\beta\,:\!\gamma\,e^{-\phi}\!:\,: \;=\; :\,:\!\beta\,\gamma\!:\,e^{-\phi}\!:\,$).

3 Subalgebra embeddings and ‘projections’ of non-critical strings

In this section we show how the embeddings of subalgebras, $s\ell(2)\hookrightarrow osp(1|2)\hookrightarrow s\ell(2|1)$, can be used to derive representations similar to (2.7)–(2.12) for the $s\ell(2)$ and $osp(1|2)$ current algebras. These representations will be in terms of the corresponding ($N\!=\!0$ and $N\!=\!1$) matter theories and a number of free fields. All the free fields originate from the $N\!=\!2$ string in the conformal gauge, and upon reduction to $N\!=\!1$ and $N\!=\!0$ they will give rise to the respective ($N\!=\!1$ or $N\!=\!0$) non-critical string ghosts and Liouville, plus an additional scalar; the $N\!=\!2$ matter sector will mix with the $N\!=\!2$ Liouville superpartners to produce the lower-supersymmetric matter theories. Thus we are going to consider the following diagram:

$$\begin{tikzcd}
\text{affine algebras} & s\ell(2) & osp(1|2) \arrow[l] & s\ell(2|1) \arrow[l] \\
\text{non-critical strings, optionally with an auxiliary scalar} & (N\!=\!0)\otimes(\partial V_*) \arrow[u] \arrow[d] & (N\!=\!1)\otimes(\partial v_*) \arrow[l] \arrow[u] \arrow[d] \arrow[dl] & N\!=\!2 \arrow[l] \arrow[u] \arrow[d] \arrow[dl] \\
\text{matter theories, optionally with Liouville fermion(s)} & (N\!=\!0)_{\rm m} \arrow[d] & (N\!=\!1)_{\rm m}\otimes(\rho) \arrow[l] \arrow[d] \arrow[dl] & (N\!=\!2)_{\rm m}\otimes(\overline{\psi},\psi) \arrow[l] \arrow[d] \arrow[dl] \\
\text{matter theories} & (N\!=\!0)_{\rm m} & (N\!=\!1)_{\rm m} & (N\!=\!2)_{\rm m}
\end{tikzcd} \tag{3.1}$$

We have adopted here the following ‘anti-mathematical’ convention as regards the direction of arrows. Consider two algebras A and B with generators (a_i) and (b_j) respectively. Then constructing the *generators* of A in terms of those of B, $a_i = f_i(b_j)$, determines what is normally denoted as the mapping $A\to B$, since it tells us that any a_i is mapped precisely into the respective f_i, viewed as an element of B. Then we would have to call, e.g., the mappings of ref. [10] *projections* rather than embeddings of string theories, like $(N\!=\!2)\to(N\!=\!1)$. We do not dare to challenge the established terminology however, and will write

$(N\!=\!1) \to (N\!=\!2)$ in the case when the $N\!=\!2$ fields are constructed out of those of an $N\!=\!1$ theory. *Then* the new mappings we are going to discuss are directed as shown in (3.1). Thus, $A \to B$ reads 'take the generators of A and construct generators of B'.

Let us begin with the embedding $osp(1|2) \hookrightarrow s\ell(2|1)$: the $osp(1|2)$ currents can be identified as

$$\begin{aligned} I^{+} &= E_{12}\,, & I^{-} &= F_{12}\,, \\ I^{+\frac{1}{2}} &= E_1 + E_2\,, & I^{-\frac{1}{2}} &= F_1 - F_2\,, \\ I^{0} &= H^{-} \end{aligned} \tag{3.2}$$

They satisfy the $osp(1|2)$ OPEs with precisely the above level k (our conventions for $osp(1|2)$ are specified in the Appendix). By substituting into (3.2) the above expressions for the $s\ell(2|1)$ currents, we will of course get a realization of the $osp(1|2)$ algebra. However that realization will not be 'irreducible' in the sense that the currents (3.2) depend actually on only certain combinations of the $N\!=\!2$ string fields. These special combinations turn out to make up an $N\!=\!1$ non-critical string theory onto which we can '*project*' from the $N\!=\!2$ string. Let us describe therefore how this is carried out. We are going to introduce a new basis in the space of fields of the $N\!=\!2$ string in such a way that some of the new fields will be the $N\!=\!1$ string fields, while the others will decouple from the string.

First, we realize the $N\!=\!2$ Liouville fermion in terms of two real fermions λ and ρ,

$$\rho(z)\,\rho(w) = -\frac{1}{z-w}\,, \qquad \lambda(z)\,\lambda(w) = \frac{1}{z-w}\,, \tag{3.3}$$

as

$$\psi = \tfrac{1}{\sqrt{2}}(\rho+\lambda)\,, \qquad \overline{\psi} = \tfrac{1}{\sqrt{2}}(\lambda-\rho)\,, \tag{3.4}$$

which, clearly, is an invertible transformation. Then, as we are going to see, ρ will be a part of the $N\!=\!1$ super-Liouville, while λ will participate in constructing an 'effective' $N\!=\!1$ matter theory: *For any* $N\!=\!2$ *matter with a non-critical central charge* $c_{\rm m} = -3-6k \neq 6$, *there is a mapping* $(N\!=\!2)_{\rm m} \otimes (\lambda) \to (N\!=\!1)_{\rm m}$ *to an* $N\!=\!1$ *theory*, which is defined by

$$\begin{aligned} G_{N=1} &= -\sqrt{2}\alpha_+ G_{\rm m} + \tfrac{1}{\sqrt{2}}(\alpha_- + \alpha_+)\overline{G}_{\rm m} + \alpha_+ H_{\rm m}\,\lambda\,, \\ T_{N=1} &= (1-\alpha_+^2)T_{\rm m} + \sqrt{2}\alpha_+^2 G_{\rm m}\,\lambda + \tfrac{1}{2}\alpha_+^2 H_{\rm m}\,H_{\rm m} + \tfrac{1}{\sqrt{2}}(\alpha_+^2 - 1)\overline{G}_{\rm m}\,\lambda + (\alpha_+^2 - \tfrac{1}{2})\partial\lambda\,\lambda\,, \end{aligned} \tag{3.5}$$

$$\alpha_+ = -\tfrac{1}{\alpha_-}\,, \qquad \alpha_-^2 = 2k+3\,. \tag{3.6}$$

The $N\!=\!1$ superconformal algebra can be checked for the generators thus constructed:

$$\begin{aligned} T_{N=1}(z)T_{N=1}(w) &= \frac{d_{N=1}/2}{(z-w)^4} + \frac{2T_{N=1}(w)}{(z-w)^2} + \frac{\partial T_{N=1}(w)}{z-w}\,, \\ T_{N=1}(z)G_{N=1}(w) &= \frac{3/2\,G_{N=1}(w)}{(z-w)^2} + \frac{\partial G_{N=1}(w)}{z-w}\,, \\ G_{N=1}(z)G_{N=1}(w) &= \frac{2/3\,d_{N=1}}{(z-w)^3} + \frac{2T_{N=1}(w)}{z-w}\,, \end{aligned} \tag{3.7}$$

the central charge being

$$d_{N=1} = \tfrac{15}{2} - 3\alpha_-^2 - 3\alpha_+^2 \tag{3.8}$$

by virtue of eqs. (A.3), (3.3). We call the theory $(G_{N=1},\, T_{N=1})$, eqs. (3.5), the 'effective' $N\!=\!1$ matter.

The remaining fields of the $N=2$ string are $\partial\overline{\phi}$, $\partial\phi$, b, c, β, γ, $\widetilde{\beta}$, $\widetilde{\gamma}$, and ξ, η. We now change the basis in the space of these free fields; as the new independent fields we take: *i)* the ghosts β, γ, b, c, *ii) the Drinfeld–Sokolov ghosts* $\widetilde{\mathsf{B}}$, $\widetilde{\Gamma}$, $\mathcal{C}$, $\mathcal{B}$, and *iii)* two real bosons ∂v_* and $\partial\varphi$, with

$$\partial v_*(z)\,\partial v_*(w)=\frac{1}{(z-w)^2}\,,\qquad \partial\varphi(z)\,\partial\varphi(w)=-\frac{1}{(z-w)^2}\,. \tag{3.9}$$

Thus, in addition to (3.4), we express the fields $\partial\overline{\phi}$, $\partial\phi$, $\widetilde{\beta}$, $\widetilde{\gamma}$, η, ξ as

$$\begin{aligned}
\partial\phi &= \alpha_+(\partial\varphi-\partial v_*)\,, & \partial\overline{\phi} &= 2\mathcal{B}\mathcal{C}+\widetilde{\mathsf{B}}\,\widetilde{\Gamma}-(\tfrac{1}{2}\alpha_--\tfrac{3}{2}\alpha_+)(\partial\varphi+\partial v_*)\,,\\
\widetilde{\beta} &= \widetilde{\mathsf{B}}\,e^{\alpha_+(\varphi-v_*)}\,, & \widetilde{\gamma} &= \widetilde{\Gamma}\,e^{-\alpha_+(\varphi-v_*)}\,,\\
\eta &= \mathcal{B}\,e^{2\alpha_+(\varphi-v_*)}\,, & \xi &= \mathcal{C}\,e^{-2\alpha_+(\varphi-v_*)}
\end{aligned} \tag{3.10}$$

The last two lines are nothing but the first and the last lines of (2.13) respectively, however with $\widetilde{\mathsf{B}}$ and $\widetilde{\Gamma}$ (bosons), and $\mathcal{C}$ and $\mathcal{B}$ (fermions) now viewed as independent fields with the operator products $\widetilde{\mathsf{B}}(z)\widetilde{\Gamma}(w)=-1/(z-w)$, $\mathcal{B}(z)\mathcal{C}(w)=1/(z-w)$. The free fermion ρ, which is what remains of the $N=2$ Liouville superpartners after λ has gone to construct the $N=1$ matter, does not mix with other fields.

In the new basis, the currents (3.2) rewrite as

$$\begin{aligned}
I^+ &= e^{2\alpha_+(\varphi-v_*)}\,,\\
I^{+\frac{1}{2}} &= \sqrt{2}\rho\,e^{\alpha_+(\varphi-v_*)}\,,\\
I^0 &= -\tfrac{3}{2}\alpha_+\partial\varphi+\tfrac{1}{2}(\alpha_-+3\alpha_+)\partial v_*+b\,c+\tfrac{1}{2}\beta\,\gamma\,,\\
I^{-\frac{1}{2}} &= \Bigl(\sqrt{2}b\,c\,\rho+\tfrac{1}{\sqrt{2}}\beta\,\gamma\,\rho+\tfrac{\alpha_-}{\sqrt{2}}\partial\varphi\,\rho+\tfrac{\alpha_-}{\sqrt{2}}G_{N=1}+\tfrac{1}{\sqrt{2}}(\alpha_-^2-2)\partial\rho\Bigr)e^{-\alpha_+(\varphi-v_*)}\,,\\
I^- &= \Bigl(-b\,c\,\beta\,\gamma-\alpha_-b\,c\,\partial\varphi+(\tfrac{3}{2}-\tfrac{1}{2}\alpha_-^2)b\,\partial c-\tfrac{1}{4}\beta\,\beta\,\gamma\,\gamma-\tfrac{1}{2}\alpha_-\beta\,\gamma\,\partial\varphi\\
&\quad+(\tfrac{1}{2}-\tfrac{1}{4}\alpha_-^2)\beta\,\partial\gamma-\tfrac{1}{4}\alpha_-^2\partial\varphi\,\partial\varphi-\tfrac{1}{2}\alpha_-\rho\,G_{N=1}+\tfrac{1}{2}\alpha_-^2T_{N=1}-(\tfrac{1}{2}+\tfrac{1}{2}\alpha_-^2)\partial b\,c\\
&\quad-\tfrac{1}{4}\alpha_-^2\partial\beta\,\gamma+\tfrac{1}{4}(\alpha_--\alpha_-^3)\partial\partial\varphi+(\tfrac{1}{4}\alpha_-^2-\tfrac{1}{2})\partial\rho\,\rho\Bigr)e^{-2\alpha_+(\varphi-v_*)}
\end{aligned} \tag{3.11}$$

and in this form represent an intrinsic-$N=1$ construction [24]: given *any* $N=1$ matter theory ($G_{N=1}$, $T_{N=1}$) determined by (3.7), (3.8), coupling it to the $N=1$ supergravity multiplet $\partial\varphi$, ρ, bc, $\beta\gamma$, and additionally tensoring with an auxiliary scalar ∂v_*, we would obtain a realization of the $osp(1|2)$ current algebra, with the $osp(1|2)$ OPEs (A.4) satisfied and the level being $k=\frac{1}{2}(\alpha_-^2-3)$.

The other two $N=2$ Drinfeld–Sokolov ghost systems BC and $\mathsf{B}\Gamma$, re-expressed in terms of the new fields introduced in (3.10),

$$\begin{aligned}
\mathsf{B} &= \beta\,e^{-\alpha_+(\varphi-v_*)}\,, & \Gamma &= \gamma\,e^{\alpha_+(\varphi-v_*)}\,,\\
B &= b\,e^{-2\alpha_+(\varphi-v_*)}\,, & C &= c\,e^{2\alpha_+(\varphi-v_*)}\,,
\end{aligned} \tag{3.12}$$

remain the ‘Drinfeld–Sokolov’ ghosts for the $osp(1|2)$ algebra: the $osp(1|2)$ theory with the twisted Sugawara energy-momentum tensor, plus the energy-momentum tensor for BC and $\mathsf{B}\Gamma$,

$$\begin{aligned}
\widetilde{T}_{osp(1|2)} &= 2\alpha_+^2(I^0\,I^0+\tfrac{1}{2}I^+\,I^-+\tfrac{1}{2}I^-\,I^++\tfrac{1}{4}I^{+\frac{1}{2}}\,I^{-\frac{1}{2}}-\tfrac{1}{4}I^{-\frac{1}{2}}\,I^{+\frac{1}{2}})+\partial I^0\,,\\
T_{\mathrm{GH}} &= -\,\partial B\,C-2B\,\partial C-\tfrac{1}{2}\partial\mathsf{B}\,\Gamma-\tfrac{3}{2}\mathsf{B}\,\partial\Gamma
\end{aligned} \tag{3.13}$$

rewrite precisely as the $N=1$ supergravity plus an extra $U(1)$ theory represented by ∂v_*:

$$\begin{aligned}
\widetilde{T}_{osp(1|2)}+T_{\mathrm{GH}} &= T_{N=1}-2b\,\partial c-\partial b\,c-\tfrac{3}{2}\beta\,\partial\gamma-\tfrac{1}{2}\partial\beta\,\gamma-\tfrac{1}{2}\partial\varphi\,\partial\varphi+\tfrac{1}{2}(\alpha_+-\alpha_-)\partial\partial\varphi-\tfrac{1}{2}\partial\rho\,\rho\\
&\quad+\tfrac{1}{2}\partial v_*\,\partial v_*+\tfrac{1}{2}(\alpha_--\alpha_+)\partial\partial v_*\,,
\end{aligned} \tag{3.14}$$

which is an $osp(1|2)$–$(N=1)$ analogue of (2.14). The central charge in the $\partial\varphi$-ρ sector is the required $\frac{15}{2}+3\alpha_-^2+3\alpha_+^2$ (cf. (3.8)).

Eqs. (3.11), (3.12) are the construction of the $osp(1|2)$ current algebra from [24], which is valid for any non-critical $N=1$ string tensored with an auxiliary scalar. Now we see that this is also the $osp(1|2)$ representation induced from the $s\ell(2|1)$ algebra realized in the $N=2$ string. The subalgebra embedding $osp(1|2)\hookrightarrow s\ell(2|1)$ thus allows us to map the respective non-critical string theories and, another level down, the matter theories as in (3.5), see (3.1).

To return to the interpretation of $(T_{N=1},G_{N=1})$ as the 'effective' $N=1$ matter made out of the ingredients of the $N=2$ string, we see that the bc and $\beta\gamma$ ghosts remain in their capacities of the reparametrization and the 'gravitino' ghosts respectively, while $\partial\varphi$ and ρ are the components of the $N=1$ super-Liouville (thus, as expected, just one of the $N=2$ supersymmetries gets broken). On the other hand, ∂v_*, $\widetilde{\mathsf{B}}\widetilde{\Gamma}$ and $\mathcal{BC}$ decouple from the $N=1$ string. It may be worth noting that the $\mathcal{BC}$ and $\widetilde{\mathsf{B}}\widetilde{\Gamma}$ ghosts make up a *topological* $bc\beta\gamma$ theory, so one may practice in proving the cohomological triviality of modding out the $\mathcal{BC}\widetilde{\mathsf{B}}\widetilde{\Gamma}$ system. The identifications of the different sectors can be summarized as follows:


$\beta\gamma$ bc $T_{\rm m}$ $H_{\rm m}$ $\overline{G}_{\rm m}$ $G_{\rm m}$ ψ $\overline{\psi}$ $\partial\phi$ $\partial\overline{\phi}$ $\widetilde{\beta}\widetilde{\gamma}$ $\xi\eta$
$\beta\gamma$ bc $T_{\rm m}$ $H_{\rm m}$ $\overline{G}_{\rm m}$ $G_{\rm m}$ λ ρ $\partial\varphi$ ∂v_* $\widetilde{\mathsf{B}}\widetilde{\Gamma}$ $\mathcal{C}\mathcal{B}$
$\beta\gamma$ bc $T_{N=1}$ $G_{N=1}$ ρ $\partial\varphi$ ∂v_* $\widetilde{\mathsf{B}}\widetilde{\Gamma}$ $\mathcal{C}\mathcal{B}$
$N=1$ string
$osp(1|2)$ $B\,C$ $\mathsf{B}\Gamma$ $\widetilde{\mathsf{B}}\widetilde{\Gamma}$ $\mathcal{C}\mathcal{B}$

We would like to stress that this picture reflects nothing but a simple fact of the subalgebra embedding $osp(1|2)\hookrightarrow s\ell(2|1)$ of the $osp(1|2)$ algebra from the bottom line into the $s\ell(2|1)$ algebra associated with the top line of the diagram.

This is not yet the end of the story: one can reduce further down to $N=0$ theories, which would correspond to taking just the (common) $s\ell(2)$ subalgebra of $s\ell(2|1)$ and $osp(1|2)$. We thus start with the theory whose energy-momentum tensor is given by the RHS of (3.14) with $\alpha_+^2\neq-1$ ($k\neq-2$) and further rearrange the fields with the aim to single out the non-critical bosonic string. First of all, the $N=1$ matter (3.7) tensored with the ρ fermion 'projects' onto the $N=0$ matter:

$$\begin{aligned}
T_{N=0} &= \frac{1}{1+\alpha_+^2}T_{N=1}-\frac{\alpha_+}{1+\alpha_+^2}G_{N=1}\,\rho+\frac{1-2\alpha_+^2}{2(1+\alpha_+^2)}\partial\rho\,\rho\\
&= \tfrac{1}{k+2}\Bigl((k+1)T_{\rm m}+G_{\rm m}\,\psi+\tfrac14 H_{\rm m}\,H_{\rm m}+\tfrac12 H_{\rm m}\,\psi\,\overline{\psi}-(k+1)\overline{G}_{\rm m}\,\overline{\psi}\\
&\qquad +\tfrac14(1+2k)\psi\,\partial\overline{\psi}-\tfrac14(1+2k)\partial\psi\,\overline{\psi})\Bigr)
\end{aligned}\tag{3.15}$$

where we have also re-expressed $T_{N=0}$ through the fields of the $N=2$ string[3]. In the following, however, we will not expand the $N=1$ fields in terms of the $N=2$ ingredients. As is easy to check, the operator products (3.7), (3.3) alone allow one to show that $T_{N=0}$ *is* an energy-momentum tensor with central charge

$$d_{N=0}=13-6(k+2)-\tfrac{6}{k+2}\,. \tag{3.16}$$

---

[3] Which shows that the throughout 'reduction' to the $N=0$ matter (and hence to the bosonic string) is possible from the $N=2$ string whenever $c_{\rm m}\neq 9$.

To 'project' onto the whole $N=0$ non-critical string, we need to identify the corresponding Liouville field. To this end we mix the free fields as follows. Recall that in the $N=1$ setting we have, in addition to the $N=1$ matter, the ghosts bc and $\beta\gamma$, the Liouville system $\partial\varphi$ and ρ, and the auxiliary scalar ∂v_*. As we saw in the first equation in (3.15), ρ goes to construct the 'effective' $N=0$ matter. Among the remaining free fields, we choose a new basis: as the new independent fields, we take the bosonic $N=1$ Drinfeld–Sokolov ghosts $\mathsf{B}\Gamma$, along with two scalars $\partial\Phi$ and ∂V_* with the operator products

$$\partial V_*(z)\,\partial V_*(w) = \frac{1}{(z-w)^2}\,, \qquad \partial\Phi(z)\,\partial\Phi(w) = -\frac{1}{(z-w)^2}\,, \tag{3.17}$$

introduced via

$$\begin{aligned}
\partial\varphi &= \alpha_+\mathsf{B}\Gamma + \tfrac{1}{\sqrt{1+\alpha_+^2}}(\partial\Phi + \alpha_+^2\partial V_*)\,,\\
\partial v_* &= \alpha_+\mathsf{B}\Gamma + \sqrt{1+\alpha_+^2}\,\partial V_*\,,\\
\beta &= \mathsf{B}\exp\Big(\tfrac{\alpha_+}{\sqrt{1+\alpha_+^2}}(\Phi - V_*)\Big)\,, \quad \gamma \;=\; \Gamma\exp\Big(\tfrac{-\alpha_+}{\sqrt{1+\alpha_+^2}}(\Phi - V_*)\Big)
\end{aligned} \tag{3.18}$$

Then the $s\ell(2)$ currents rewrite as

$$\begin{aligned}
I^+ &= e^{\sqrt{2}a_+(\Phi - V_*)}\\
I^0 &= -\sqrt{2}a_+\partial\Phi + (\tfrac{1}{\sqrt{2}}a_- + \sqrt{2}a_+)\partial V_* + bc\\
I^- &= \Big((2-a_-^2)b\partial c - \sqrt{2}a_-\partial\Phi bc - a_-^2\partial bc\\
&\qquad + a_-^2\big(T_{N=0} - \tfrac{1}{2}\partial\Phi\partial\Phi - \tfrac{1}{\sqrt{2}}(a_- + a_+)\partial\partial\Phi\big)\Big)e^{-\sqrt{2}a_+(\Phi - V_*)}
\end{aligned} \tag{3.19}$$

where $a_- = -1/a_+$ and

$$a_+ = \frac{\sqrt{2}\alpha_+}{\sqrt{1+\alpha_+^2}}\,, \tag{3.20}$$

whence $a_-^2 = k+2$. The $s\ell(2)$ currents are thus expressed through the energy-momentum tensor $T_{N=0}$, the bc (reparametrization) ghosts, the $N=0$ Liouville $\partial\Phi$ and the auxiliary scalar ∂V_*. In fact, any $N=0$ matter theory with central charge (3.16) can be dressed into an $s\ell(2)$ algebra of level k by tensoring with the ghosts, a 'Liouville' scalar and an auxiliary scalar. This construction, known from [23], is therefore recovered from the embedding $s\ell(2) \hookrightarrow osp(1|2)$ [24].

The fermionic Drinfeld–Sokolov ghosts BC that we had in the $N=1$ context, remain the Drinfeld–Sokolov ghosts for the $s\ell(2)$ algebra. They are re-expressed in terms of the new ingredients as

$$B = b\,e^{-\sqrt{2}a_+(\Phi - V_*)}\,, \qquad C = c\,e^{\sqrt{2}a_+(\Phi - V_*)}\,, \tag{3.21}$$

which allows us to show that the (twisted) $s\ell(2)$ Sugawara energy-momentum tensor, plus the energy-momentum tensor for the BC ghosts, evaluate as the $N=0$ string with an auxiliary ∂V_* scalar:

$$\begin{aligned}
&\tfrac{1}{k+2}(I^0\,I^0 + \tfrac{1}{2}I^+\,I^- + \tfrac{1}{2}I^-\,I^+) + \partial I^0 - \partial B\,C - 2B\,\partial C\\
&\qquad = T_{N=0} - 2b\,\partial c - \partial b\,c - \tfrac{1}{2}\partial\Phi\,\partial\Phi - \tfrac{1}{\sqrt{2}}(a_- - a_+)\partial\partial\Phi\\
&\qquad + \tfrac{1}{2}\partial V_*\,\partial V_* + \tfrac{1}{\sqrt{2}}(a_- - a_+)\partial\partial V_*\,.
\end{aligned} \tag{3.22}$$

This is the $N=0$ version of the 'completeness relation'. The central charge in the $\partial\Phi$ ('Liouville') sector is $13 + 6(k+2) + \frac{6}{k+2}$, as it should.

A remark is in order concerning the mapping ($N=1$ matter, ρ) → (‘effective’ $N=0$ matter), eq. (3.15), induced by the embedding $s\ell(2) \hookrightarrow osp(1|2)$: it relates the central charges as

$$d_{N=1} = \frac{d_{N=0}^2 + 13d_{N=0} - 5 \pm (d_{N=0} - 1)\sqrt{(25 - d_{N=0})(1 - d_{N=0})}}{2(2 + d_{N=0})} \tag{3.23}$$

When $d_{N=0} \to -2$, the ‘+’-branch gives a finite value $d_{N=1} = -\frac{5}{2}$. The inverse relation reads

$$d_{N=0} = \frac{(d_{N=1} - \frac{27}{2})(d_{N=1} + \frac{1}{2}) \pm (\frac{3}{2} - d_{N=1})\sqrt{(\frac{27}{2} - d_{N=1})(\frac{3}{2} - d_{N=1})}}{2d_{N=1} - 27} \tag{3.24}$$

The mapping (3.24) can be applied to any $N=1$ (p,q) minimal model tensored with a ρ fermion. The $N=1$ central charge is then mapped as

$$\tfrac{3}{2} - 3\tfrac{p}{q} - 3\tfrac{q}{p} \mapsto \begin{cases} 13 - 6\frac{p+q}{2q} - 6\frac{2q}{p+q} \\ 13 - 6\frac{p+q}{2p} - 6\frac{2p}{p+q} \end{cases} \tag{3.25}$$

depending on the sign in (3.24), and the ‘effective’ $N=0$ matter theory thus gets identified with a model from the $N=0$ minimal series.

The main feature that distinguishes the $s\ell(2|1)/N=2$ case is that, in contrast to $osp(1|2)$ and $s\ell(2)$, the construction of the $s\ell(2|1)$ currents does *not* involve any extra free fields in addition to the fields of the respective supergravity in the conformal gauge. Note also that the representation (3.19) of $s\ell(2)$ in terms of the $N=0$ non-critical string $(T_{N=0}, \partial\Phi, b, c)$ tensored with the ∂V_* scalar blows up at $k \to -2$, which naturally corresponds to the infinite matter central charge $d_{N=0} \to \infty$; similarly, the construction (3.11) of $osp(1|2)$ works only for $k \neq -\frac{3}{2}$, which means having finite $N=1$ matter central charges $d_{N=1} \neq \infty$. On the other hand, the realization (2.7)–(2.12) of the $s\ell(2|1)$ algebra remains valid for any finite level k.

In the *critical* $N=2$ string, with $c_{\rm m} = 6$, $q^2 = -\frac{1}{2}$, the Liouville sector decouples from the string, but not from the $s\ell(2|1)$ algebra, which continues smoothly to the corresponding level $k = -\frac{3}{2}$. The construction (2.7)–(2.12) is therefore somewhat ‘less natural’, from the stringy point of view, for the critical $N=2$ string, since it requires either introducing an ‘external’ $\partial\phi\partial\overline{\phi}\,\psi\overline{\psi}$-theory or structuring the critical matter into the ‘proper matter’ and the ‘Liouville’ theory whose central charges add up to the critical value.

4 Topological strings

In this section we reformulate eqs. (2.7)–(2.12) for the case of the *topological* strings [26]. We thus start with the topological conformal matter represented by the generators $\mathcal{T}_{\rm m}$, $\mathcal{G}_{\rm m}$, $\mathcal{Q}_{\rm m}$, and $\mathcal{H}_{\rm m}$, which satisfy the twisted-$N=2$ algebra

$$\begin{aligned}
\mathcal{T}_{\rm m}(z)\mathcal{T}_{\rm m}(w) &= \frac{2\mathcal{T}_{\rm m}(w)}{(z-w)^2} + \frac{\partial\mathcal{T}_{\rm m}}{z-w}\,, & \mathcal{H}_{\rm m}(z)\mathcal{H}_{\rm m}(w) &= \frac{\mathsf{c}/3}{(z-w)^2}\,,\\
\mathcal{T}_{\rm m}(z)\mathcal{G}_{\rm m}(w) &= \frac{2\mathcal{G}_{\rm m}(w)}{(z-w)^2} + \frac{\partial\mathcal{G}_{\rm m}}{z-w}\,, & \mathcal{T}_{\rm m}(z)\mathcal{Q}_{\rm m}(w) &= \frac{\mathcal{Q}_{\rm m}(w)}{(z-w)^2} + \frac{\partial\mathcal{Q}_{\rm m}}{z-w}\,,\\
\mathcal{T}_{\rm m}(z)\mathcal{H}_{\rm m}(w) &= \frac{\mathsf{c}/3}{(z-w)^3} + \frac{\mathcal{H}_{\rm m}(w)}{(z-w)^2} + \frac{\partial\mathcal{H}_{\rm m}}{z-w}\,, & &\\
\mathcal{H}_{\rm m}(z)\mathcal{G}_{\rm m}(w) &= \frac{\mathcal{G}_{\rm m}}{z-w}\,, & \mathcal{H}_{\rm m}(z)\mathcal{Q}_{\rm m}(w) &= -\frac{\mathcal{Q}_{\rm m}}{z-w}\,,\\
\mathcal{Q}_{\rm m}(z)\mathcal{G}_{\rm m}(w) &= \frac{\mathsf{c}/3}{(z-w)^3} - \frac{\mathcal{H}_{\rm m}(w)}{(z-w)^2} + \frac{\mathcal{T}_{\rm m}}{z-w}\,, & &
\end{aligned} \tag{4.1}$$

with the topological central charge

$$\mathsf{c} = -3 - 6k\,. \tag{4.2}$$

We will not introduce more notations and will have $\overline{\psi}$ denote the gradient of the 'topological' fermion χ, for which $\partial\chi(z)\psi(w) = 1/(z-w)$. Similarly, we will preserve the notation $\partial\overline{\phi}$ for what is often called $\partial\pi$ in the topological string [26]. The ghost sector will consist of the bosonic and fermionic ghosts β, γ and b, c [26]. We thus have the following fields making up the topological string, with their conformal dimensions given in Table 1.

b	c	β	γ	$\mathcal{T}_{\rm m}$	$\mathcal{G}_{\rm m}$	$\mathcal{Q}_{\rm m}$	$\mathcal{H}_{\rm m}$	$\partial\overline{\phi}$	$\partial\phi$	ψ	$\overline{\psi}$
2	-1	2	-1	2	2	1	1	1	1	0	1

Table 1. Conformal dimensions of the topological string ingredients

Then the formulae (2.8)–(2.12) can very easily [4] be modified to produce the $s\ell(2|1)$ algebra constructed out of the fields from Table 1:

$$\begin{aligned}
\mathcal{H}^+ &= -\tfrac{1}{2}\mathcal{H}_{\rm m} + \tfrac{1}{2}\psi\,\overline{\psi}\,,\\
\mathcal{H}^- &= -\tfrac{1}{2}\partial\overline{\phi} + \tfrac{1}{2}(-3+k)\partial\phi + \tfrac{1}{2}(\tfrac{9}{2\mathsf{q}^2} + \tfrac{\mathsf{q}^2}{3})b\,c + \tfrac{1}{2}(\tfrac{9}{2\mathsf{q}^2} - \tfrac{\mathsf{q}^2}{3})\beta\,\gamma\,,\\
\mathcal{F}_1 &= \Big(\mathcal{G}_{\rm m} - \tfrac{1}{2}(\tfrac{9}{2\mathsf{q}^2} + \tfrac{\mathsf{q}^2}{3})bc\,\overline{\psi} - \tfrac{1}{2}(\tfrac{9}{2\mathsf{q}^2} - \tfrac{\mathsf{q}^2}{3})\beta\gamma\,\overline{\psi} + \tfrac{1}{2}\partial\overline{\phi}\,\overline{\psi}\\
&\quad + \tfrac{1}{2}(3+k)\partial\phi\,\overline{\psi} - \tfrac{1}{2}\mathcal{H}_{\rm m}\overline{\psi} - (k+\tfrac{1}{2})\partial\overline{\psi}\Big)e^{-\phi}\,,\\
\mathcal{F}_2 &= \Big((k+1)\mathcal{Q}_{\rm m} + \tfrac{1}{2}(\tfrac{9}{2\mathsf{q}^2} + \tfrac{\mathsf{q}^2}{3})b\,c\,\psi + \tfrac{1}{2}(\tfrac{9}{2\mathsf{q}^2} - \tfrac{\mathsf{q}^2}{3})\beta\,\gamma\,\psi - \tfrac{1}{2}\partial\overline{\phi}\,\psi\\
&\quad - \tfrac{1}{2}(3+k)\partial\phi\,\psi - \tfrac{1}{2}\mathcal{H}_{\rm m}\,\psi + (k+\tfrac{1}{2})\partial\psi\Big)e^{-\phi}\,,
\end{aligned} \tag{4.3}$$

while $\mathcal{E}_i = E_i$, eqs. (2.7) (and $\mathcal{F}_{12}$ uniquely determined by $\mathcal{F}_1$ and $\mathcal{F}_2$). Here q is a free parameter, which will be related to the topological Liouville background charge. To see how this can be done, consider the 'topological' analogue of the 'completeness relation' (2.14). The formulae (4.3) differ from (2.8)–(2.11) essentially by a redefinition of the fields by means of an $O(2,2)$ rotation in the space of the four ghost currents bc, $\beta\gamma$, $\widetilde{\beta}\widetilde{\gamma}$, and $\eta\xi$: the bc and $\beta\gamma$ currents in (4.3) are particular linear combinations of the four currents from (2.8)–(2.11). Then the construction of the 'Drinfeld–Sokolov' ghosts changes accordingly: in the topological setting, we define, instead of (2.13),

$$\begin{aligned}
\mathsf{B}_{\rm t} &= \beta\,e^{-(9/(2\mathsf{q}^2)-\mathsf{q}^2/3)\phi}\,, & \Gamma_{\rm t} &= \gamma\,e^{(9/(2\mathsf{q}^2)-\mathsf{q}^2/3)\phi}\,,\\
B_{\rm t} &= b\,e^{-(9/(2\mathsf{q}^2)+\mathsf{q}^2/3)\phi}\,, & C_{\rm t} &= c\,e^{(9/(2\mathsf{q}^2)+\mathsf{q}^2/3)\phi}\,.
\end{aligned} \tag{4.4}$$

Now, we evaluate the energy-momentum tensor for these 'Drinfeld–Sokolov' [5] ghosts and add it with the $s\ell(2|1)$ Sugawara energy-momentum tensor; the latter, however, has to be *doubly twisted*, then it follows from the previous formulae that

$$\begin{aligned}
&\tfrac{1}{k+1}\Big(\mathcal{H}^-\mathcal{H}^- - \mathcal{H}^+\mathcal{H}^+ + \tfrac{1}{2}\mathcal{E}_{12}\,\mathcal{F}_{12} + \tfrac{1}{2}\mathcal{F}_{12}\,\mathcal{E}_{12} + \tfrac{1}{2}\mathcal{E}_1\,\mathcal{F}_1 - \tfrac{1}{2}\mathcal{F}_1\,\mathcal{E}_1 - \tfrac{1}{2}\mathcal{E}_2\,\mathcal{F}_2 + \tfrac{1}{2}\mathcal{F}_2\,\mathcal{E}_2\Big) + \partial\mathcal{H}^- + \partial\mathcal{H}^+\\
&\qquad - \partial B_{\rm t}\,C_{\rm t} - 2B_{\rm t}\,\partial C_{\rm t} - \partial\mathsf{B}_{\rm t}\,\Gamma_{\rm t} - 2\mathsf{B}_{\rm t}\,\partial\Gamma_{\rm t}\\
&\qquad\qquad = \mathcal{T}_{\rm m} - \partial b\,c - 2b\,\partial c - \partial\beta\,\gamma - 2\beta\,\partial\gamma - \partial\overline{\phi}\,\partial\phi + \partial\psi\,\overline{\psi} + (\mathsf{q}^2 + k)\partial\partial\phi\,,
\end{aligned} \tag{4.5}$$

---

[4] The reason being that F_1, F_2 and H^- (and hence F_{12}) depend only on the combination $\beta\,\gamma + \widetilde{\beta}\,\widetilde{\gamma} + 2\eta\,\xi + 2bc$, as can be seen from (2.16)–(2.17).

[5] In the topological context, the name may be misleading, but we continue to use it in order to keep track of the various ghost theories.

thereby recovering (for $\overline{\psi}=\partial\chi$) the energy-momentum tensor of the topological string.

Dimensions of the 'composite' fields evaluated with respect to the energy-momentum tensor on the LHS of (4.5), are as follows:

$\mathcal{E}_1$	$\mathcal{E}_2$	$\mathcal{E}_{12}$	$\mathcal{H}^+$	$\mathcal{H}^-$	$\mathcal{F}_1$	$\mathcal{F}_2$	$\mathcal{F}_{12}$	B_{t}	Γ_{t}	B_{t}	C_{t}
0	1	0	1	1	2	1	2	2	-1	2	-1

Table 2. Conformal dimensions of doubly-twisted $s\ell(2|1)$ currents and Drinfeld–Sokolov ghosts

5 Constructing the highest-weight states and a KPZ-like picture

In this section, we go further with the analysis of the new representation for the $s\ell(2|1)$ algebra and consider briefly how states in the representation space of $s\ell(2|1)$ can be built in terms of the 'stringy' ingredients. We begin with the $s\ell(2|1)$ construction induced from the topological string, since this would allow us to have fewer fields without missing any general feature.

In the ghost sectors, we choose the $sl(2)$-invariant vacua [41]

$$b_{\geq -1}|0\rangle_{bc}=0\,,\quad c_{\geq 2}|0\rangle_{bc}=0\,,\qquad \beta_{\geq -1}|0\rangle_{\beta\gamma}=0\,,\quad \gamma_{\geq 2}|0\rangle_{\beta\gamma}=0\,. \tag{5.1}$$

In the $\overline{\psi}\psi$-sector, similarly,

$$\overline{\psi}_{\geq 0}|0\rangle_{\overline{\psi}\psi}=0\,,\quad \psi_{\geq 1}|0\rangle_{\overline{\psi}\psi}=0\,. \tag{5.2}$$

Consider further the topological matter sector (4.1). There are two different types of highest-weight conditions for the $N=2$ algebra: *i)* those that define 'chiral' primary states [32],

$$\begin{aligned}
\mathcal{Q}_{\geq 1}\,|h,\ell\rangle &= \mathcal{G}_{\geq 0}\,|h,\ell\rangle \;=\; \mathcal{L}_{\geq 1}\,|h,\ell\rangle \;=\; \mathcal{H}_{\geq 1}\,|h,\ell\rangle \;=\; 0\,,\\
\mathcal{H}_0\,|h\rangle &= h\,|h,\ell\rangle\,,\\
\mathcal{L}_0\,|h\rangle &= \ell\,|h,\ell\rangle
\end{aligned} \tag{5.3}$$

(where $\mathcal{T}(z)=\sum\mathcal{L}_n\,z^{-n-2}$, $\mathcal{G}(z)=\sum\mathcal{G}_n\,z^{-n-2}$, $\mathcal{Q}(z)=\sum\mathcal{Q}_n\,z^{-n-1}$, $\mathcal{H}(z)=\sum\mathcal{H}_n\,z^{-n-1}$), and *ii)* those that define 'chiral' and at the same time 'BRST-invariant' primary states,

$$\begin{aligned}
\mathcal{Q}_{\geq 0}\,|h\rangle &= \mathcal{G}_{\geq 0}\,|h\rangle \;=\; \mathcal{L}_{\geq 1}\,|h\rangle \;=\; \mathcal{H}_{\geq 1}\,|h\rangle \;=\; 0\,,\\
\mathcal{H}_0\,|h\rangle &= h\,|h\rangle\,,\\
\mathcal{L}_0\,|h\rangle &= 0\,.
\end{aligned} \tag{5.4}$$

In [21], the latter were called *topological* primary states, but the term may be a bit overloaded in the present paper.

The $s\ell(2|1)$ highest-weight states can be obtained by tensoring the corresponding $N=2$ highest-weight states with the free-field vacua. The parameters have to be related in order that the $s\ell(2|1)$ state satisfy the required conditions. In this paper we only consider how this works for the $N=2$ topological highest-weight states. In fact, the $s\ell(2|1)$ highest-weight states come out tensored with the 'Drinfeld–Sokolov' ghost vacua: consider

$$|p,h\rangle_{s\ell(2|1)}\otimes|0\rangle_{B_{\mathrm{t}}C_{\mathrm{t}}}\otimes|0\rangle_{\mathsf{B}_{\mathrm{t}}\Gamma_{\mathrm{t}}}=|0\rangle_{bc}\otimes|0\rangle_{\beta\gamma}\otimes|0\rangle_{\overline{\psi}\psi}\otimes|h\rangle\otimes|p\rangle_{\overline{\phi}\phi} \tag{5.5}$$

where $|p\rangle_{\overline{\phi}\phi}$ is a state corresponding to the vertex operator $\exp p\,\phi$. From (4.4) it follows that (5.5) *is* a proper vacuum for each of the 'Drinfeld–Sokolov' ghost systems. For the $s\ell(2|1)$ currents (4.3), similarly, we find (with the mode expansions of the currents performed in accordance with their dimensions from Table 2):

$$\begin{array}{rclrclrcl}
(\mathcal{E}_1)_{\geq 1}|p,h\rangle_{s\ell(2|1)} &=& 0\,, & (\mathcal{E}_2)_{\geq 0}|p,h\rangle_{s\ell(2|1)} &=& 0\,, & (\mathcal{E}_{12})_{\geq 1}|p,h\rangle_{s\ell(2|1)} &=& 0\,,\\
(\mathcal{F}_1)_{\geq 0}|p,h\rangle_{s\ell(2|1)} &=& 0\,, & (\mathcal{F}_2)_{\geq 1}|p,h\rangle_{s\ell(2|1)} &=& 0\,, & (\mathcal{F}_{12})_{\geq 1}|p,h\rangle_{s\ell(2|1)} &=& 0\,,\\
\mathcal{H}^-_{\geq 1}|p,h\rangle_{s\ell(2|1)} &=& 0\,, & \mathcal{H}^+_{\geq 1}|p,h\rangle_{s\ell(2|1)} &=& 0\,, &&&
\end{array} \tag{5.6}$$

In addition, we find the following eigenvalues of the Cartan operators:

$$\begin{array}{rcl}
\mathcal{H}^+_0\,|p,h\rangle_{s\ell(2|1)} &=& -\frac{h}{2}\,|p,h\rangle_{s\ell(2|1)}\,,\\
\mathcal{H}^-_0\,|p,h\rangle_{s\ell(2|1)} &=& \frac{p}{2}\,|p,h\rangle_{s\ell(2|1)}\,,
\end{array} \tag{5.7}$$

which actually label the states we are considering[6].

Those modes of the currents that do not annihilate the state thus constructed, do generate a module from $|p,h\rangle_{s\ell(2|1)}$. In terms of the 'component' fields from Table 1, these are evaluated e.g. as

$$\begin{array}{rcl}
(\mathcal{F}_1)_{-1}|p,h\rangle_{s\ell(2|1)} &=& |0\rangle_{bc}\otimes|0\rangle_{\beta\gamma}\otimes|0\rangle_{\overline{\psi}\psi}\otimes(\mathcal{G}_{\rm m})_{-1}|h\rangle\otimes|p\rangle_{\overline{\phi}\phi}\\
&& -\,\frac{h+p}{2}\,|0\rangle_{bc}\otimes|0\rangle_{\beta\gamma}\otimes\overline{\psi}_{-1}|0\rangle_{\overline{\psi}\psi}\otimes|h\rangle\otimes|p\rangle_{\overline{\phi}\phi}\,,\\
(\mathcal{F}_2)_0|p,h\rangle_{s\ell(2|1)} &=& \frac{p-h}{2}\,|0\rangle_{bc}\otimes|0\rangle_{\beta\gamma}\otimes\psi_0|0\rangle_{\overline{\psi}\psi}\otimes|h\rangle\otimes|p\rangle_{\overline{\phi}\phi}\,,
\end{array} \tag{5.8}$$

etc.

However, the absence of the annihilation condition for $(\mathcal{F}_{12})_0$ (which would be $(F_{12})_1$ in the standard setting with the currents being of dimension 1, as corresponds to the *un*twisted Sugawara energy-momentum tensor) means that we are in the $s\ell(2|1)$ module with infinitely many highest-weight vectors (thus, in a sense 'without' a highest-weight vector, assuming this to be unique); our aim, on the other hand, is to build an $s\ell(2|1)$ Verma module with a unique highest-weight state out of the ingredients of the $N=2$ string. This requires introducing further relations between the parameters involved, which entails the restriction to a class of 'overdetermined' $s\ell(2|1)$ highest-weight states. An obvious possibility is to set $p=h$, then we see from (5.8) that $(\mathcal{F}_2)_0$ would vanish on the highest-weight state, and *hence* so would do $(\mathcal{F}_{12})_0$.

For the honest $N=2$ string, such a mechanism of having, in that case, $(F_2)_0$ and $(F_{12})_1$ vanish simultaneously on the highest-weight state works as follows. First of all, the construction similar to (5.5) follows by tensoring both sides of (5.5) with the missing ghosts and then performing the appropriate $O(2,2)$ mixing in the ghost sector. Then we would have $(E_1)_{\geq 0}\approx 0$, $(E_2)_{\geq 0}\approx 0$, $(E_{12})_{\geq 0}\approx 0$, $(F_1)_{\geq 1}\approx 0$, $(F_2)_{\geq 1}\approx 0$, $(F_{12})_{\geq 2}\approx 0$ on the thus constructed state. Further, the energy-momentum tensor (2.5) endows it with dimension $(p-h)/2$. Now, requiring this to vanish will also result in the vanishing of the total $U(1)$ charge[7] and, on the other hand, *will also be the condition for* $(F_2)_0$ *and* $(F_{12})_1$ *to annihilate the highest-weight state* (with the other highest-weight conditions already satisfied). Thus the requirement that the free fields dress the $N=2$ topological highest-weight state into a *massless* $s\ell(2|1)$ state produces the restricted class of the $s\ell(2|1)$ highest-weight conditions.

Thus, the $N=2$ string appears to be a 'self-KPZ' theory, for which the appropriate generalizations of the DDK ansatz [27, 28] in the conformal gauge and the KPZ formulation [29, 30] can be carried out on the same field space.

---

[6] The third parameter, the eigenvalue k of the central element of the algebra, is omitted from the notations.

[7] To avoid confusion, note that our normalization of the Liouville currents $\partial\phi$ and $\partial\overline{\phi}$ differs from that of ref. [17] by a transformation of the form (2.4).

We would like to stress that one has the *Verma* module over the highest-weight vector (5.4) in the $N=2$ matter sector (hence, in particular, no 'accidental' vanishing of singular vectors can occur unless one bosonizes the matter sector through free fields, thereby going over to Fock modules).

6 A 'non-stringy' reformulation and the Wakimoto representations of the affine $s\ell(2|1)$

Our construction of section 2 employed the full set of fields of the $N=2$ string in the conformal gauge. As a result, we were able to construct not only the $s\ell(2|1)$ currents, but also the Drinfeld–Sokolov ghosts; the theory $s\ell(2|1)\otimes$ (DS ghosts) was singled out by the 'completeness relation' (2.14), whose RHS gives the $N=2$ string energy-momentum tensor. However, when one is not particularly interested in the string, one may prefer having fewer ingredients at the starting point and, accordingly, only the $s\ell(2|1)$ currents (with no extra 'Drinfeld–Sokolov' ghosts) as the output. Such a reformulation is easy to arrive at, by a change of basis in the space of the free fields: in addition to (2.16), one introduces

$$\partial U = -\tfrac{1}{2}\partial\overline{\phi} - \tfrac{1}{2}(3-k)\partial\phi + b\,c + \tfrac{1}{2}\beta\,\gamma + \tfrac{1}{2}\widetilde{\beta}\,\widetilde{\gamma} + \eta\,\xi \tag{6.1}$$

Then ∂F and ∂U, with

$$\partial F(z)\partial F(w) = \frac{-k/2}{(z-w)^2}\,, \qquad \partial U(z)\partial U(w) = \frac{k/2}{(z-w)^2}\,, \tag{6.2}$$

can be considered as new independent free fields, and the $s\ell(2|1)$ currents at level $k\neq 0$ rewrite in terms of $T_{\rm m}$, $H_{\rm m}$, $G_{\rm m}$, $\overline{G}_{\rm m}$, F, U, ψ, and $\overline{\psi}$ as

$$\begin{aligned}
E_1 &= \psi\, e^{\frac{1}{k}(U-F)}\,, \qquad E_2 = \overline{\psi}\, e^{\frac{1}{k}(U-F)}\,, \qquad E_{12} = e^{\frac{2}{k}(U-F)}\,,\\
H^+ &= -\tfrac{1}{2}H_{\rm m} + \tfrac{1}{2}\psi\,\overline{\psi}\,, \qquad H^- = \partial U\,,\\
F_1 &= (G_{\rm m} - \overline{\psi}\,\partial F - \tfrac{1}{2}H_{\rm m}\,\overline{\psi} - (k+\tfrac{1}{2})\partial\overline{\psi})e^{-\frac{1}{k}(U-F)}\,,\\
F_2 &= ((k+1)\overline{G}_{\rm m} + \psi\,\partial F - \tfrac{1}{2}H_{\rm m}\,\psi + (k+\tfrac{1}{2})\partial\psi)e^{-\frac{1}{k}(U-F)}\,,\\
F_{12} &= (-\partial F\,\partial F - (k+1)\partial\partial F + (k+2)T_{N=0})e^{-\frac{2}{k}(U-F)}\,,
\end{aligned} \tag{6.3}$$

where the 'effective' energy-momentum tensor $T_{N=0}$ stands for the second expression in (3.15). Equations (6.3) show, in particular, that ∂U is nothing but a 'bosonization' of the diagonal current from the $s\ell(2)$ subalgebra[8]; thus, factoring it away would leave us with the appropriate super-parafermions, which, too, would be represented in terms of an arbitrary $N=2$ matter with the appropriate central charge and the free fields F, $\overline{\psi}$, ψ. The Sugawara energy-momentum tensor (A.2) evaluates in the realization (6.3) as

$$T_{\rm Sug} = T_{\rm m} - \tfrac{1}{k}\partial F\,\partial F - \partial\partial F - \tfrac{1}{2}\psi\,\partial\overline{\psi} + \tfrac{1}{2}\partial\psi\,\overline{\psi} + \tfrac{1}{k}\partial U\,\partial U \tag{6.4}$$

The principle that (6.3) inherits from the construction of section 2 is that the $N=2$ matter sector can be arbitrary.

Recently, the *Wakimoto* $s\ell(2|1)$ modules [25] and their relation to the $N=2$ strings were considered in [34]. As was pointed out in [23] on the $s\ell(2)$ example, representations of the type of those considered in the present paper (in particular, eqs. (6.3)) can be mapped onto the Wakimoto representations at the expense of explicitly bosonizing the matter sector through some free fields. The same is true for the $s\ell(2|1)$ algebra, however there are some new features due to the existence of two simple root systems not related via a proper affine Weyl group reflection.

---

[8] One should probably have normalized the ∂F and ∂U currents to ± 1 over the poles in (6.2), in which case the exponents in (6.3) would acquire the factors $\pm\sqrt{\frac{1}{2k}}$, $\pm\sqrt{\frac{2}{k}}$, and the energy-momentum tensor in (6.4) would take the canonical form. An essential point is that, in whatever normalization, the currents ∂U and ∂F have opposite signatures.

The fact that $s\ell(2|1)$ admits two different simple root systems manifests itself in the hamiltonian reduction: there exist two different ways to perform the hamiltonian reduction of $s\ell(2|1)$ to $N=2$ matter [16]. Another consequence of the existence of two simple root systems has been worked out in [34], where a 'second' Wakimoto representation for $s\ell(2|1)$ was found, related to the appropriate 'second' Gauss decomposition. An interesting question thus is whether the two essentially different Wakimoto representations are related in any reasonable way to the realization (6.3). As we have noted, the $s\ell(2)$ experience tells us that the Wakimoto ingredients can indeed be constructed as soon as one explicitly 'bosonizes' the matter sector in terms of free fields. Such a bosonization is just what we have been carefully avoiding, since an important property of the realization (6.3) is that it is valid for an *arbitrary* matter sector. Now, in contrast to our strategy so far, we do bosonize the $N=2$ matter sector ($T_{\rm m}$, $G_{\rm m}$, $H_{\rm m}$, $\overline{G}_{\rm m}$) and see if the necessary ingredients of the Wakimoto representation can be constructed in the resulting space of free fields.

However, there are *two* essentially different ways to 'bosonize' the $N=2$ matter. The first one is through a complex scalar and a fermion,

$$\partial\overline{X}(z)\,\partial X(w)=\frac{1}{(z-w)^2}\,,\qquad \chi(z)\,\overline{\chi}(w)=-\frac{1}{z-w}\,,\tag{6.5}$$

as in [35, 36, 37] (modulo simple field redefinitions):

$$\begin{aligned}
T_{\rm m} &= \tfrac{1}{2}\chi\,\partial\overline{\chi}-\tfrac{1}{2}\partial\chi\,\overline{\chi}+\partial\overline{X}\,\partial X-\tfrac{1}{2}\partial\partial\overline{X}-\tfrac{1}{2}(k+1)\partial\partial X\,,\\
H_{\rm m} &= \partial\overline{X}-(k+1)\partial X-\chi\,\overline{\chi}\,,\\
\overline{G}_{\rm m} &= -\overline{\chi}\,\partial\overline{X}+(k+1)\partial\overline{\chi}\,,\\
G_{\rm m} &= \chi\,\partial X-\partial\chi\,.
\end{aligned}\tag{6.6}$$

The operator products (6.5) are invariant under rescalings similar to (2.6), and this freedom, both in the $\partial\overline{X}\partial X$ and $\overline{\chi}\chi$ sectors, has been used in order to minimize the number of times $(k+1)^{-1}$ appears in (6.6) and the subsequent formulae.

The following formulae show why we needed the fields χ, $\overline{\chi}$, $\partial\overline{X}$ and ∂X to represent the $N=2$ matter: the Wakimoto ingredients are constructed in terms of the free fields from (6.3) and (6.6) as

$$\begin{aligned}
\beta_{\rm w} &= e^{\frac{2}{k}(U-F)}\,,\\
\gamma_{\rm w} &= \Bigl(\partial F+\tfrac{1}{2}(k+1)\partial X+\tfrac{1}{2}\partial\overline{X}+\tfrac{1}{2}\psi\,\overline{\psi}-\tfrac{1}{2}\chi\,\overline{\chi}-(k+1)\overline{\psi}\,\overline{\chi}\Bigr)e^{-\frac{2}{k}(U-F)}\,,\\
\partial\overline{\varphi}_{\rm w} &= \partial\overline{X}-\tfrac{k+1}{k}\partial U+\tfrac{k+1}{k}\partial F\,, & \partial\varphi_{\rm w} &= \partial X+\tfrac{1}{k}\partial F-\tfrac{1}{k}\partial U\,,\\
\overline{\Psi}^{(1)}_{\rm w} &= \chi\,e^{-\frac{1}{k}(U-F)}+(k+1)\overline{\psi}\,e^{-\frac{1}{k}(U-F)}\,, & \Psi^{(1)}_{\rm w} &= \overline{\chi}\,e^{\frac{1}{k}(U-F)}\,,\\
\overline{\Psi}^{(2)}_{\rm w} &= \overline{\psi}\,e^{\frac{1}{k}(U-F)}\,, & \Psi^{(2)}_{\rm w} &= \psi\,e^{-\frac{1}{k}(U-F)}+(k+1)\overline{\chi}\,e^{-\frac{1}{k}(U-F)}
\end{aligned}\tag{6.7}$$

which would be impossible had we had only the generators $T_{\rm m}$, $G_{\rm m}$, $\overline{G}_{\rm m}$ and $H_{\rm m}$ (and the remaining fields F, U, and $\psi,\overline{\psi}$). The Wakimoto fields (6.7) can be considered as an independent bosonic beta-gamma system, a complex boson and two fermionic bc systems with operator products

$$\begin{aligned}\beta_{\mathrm{W}}(z)\,\gamma_{\mathrm{W}}(w) &= \frac{-1}{z-w}\,, & \partial\varphi_{\mathrm{W}}(z)\,\partial\overline{\varphi}_{\mathrm{W}}(w) &= \frac{1}{(z-w)^2}\,,\\ \Psi^{(1)}_{\mathrm{W}}(z)\,\overline{\Psi}^{(1)}_{\mathrm{W}}(w) &= \frac{-1}{z-w}\,, & \Psi^{(2)}_{\mathrm{W}}(z)\,\overline{\Psi}^{(2)}_{\mathrm{W}}(w) &= \frac{1}{z-w}\,.\end{aligned} \tag{6.8}$$

Now we insert the fields (6.7) into the Wakimoto realization for $s\ell(2|1)$. A convenient form of the latter is obtained from the formulae of refs. [25, 34] by simple field redefinitions which are automorphisms of the defining operator product expansions:

$$\begin{aligned}E_1 &= \Psi^{(2)}_{\mathrm{W}}\,\beta_{\mathrm{W}} - (k+1)\Psi^{(1)}_{\mathrm{W}}\,, \qquad E_2 = \overline{\Psi}^{(2)}_{\mathrm{W}}\,, \qquad E_{12} = \beta_{\mathrm{W}}\,,\\ H^+ &= -\tfrac{1}{2}\partial\overline{\varphi}_{\mathrm{W}} + \tfrac{1}{2}(k+1)\partial\varphi_{\mathrm{W}} - \tfrac{1}{2}\Psi^{(1)}_{\mathrm{W}}\,\overline{\Psi}^{(1)}_{\mathrm{W}} + \tfrac{1}{2}\Psi^{(2)}_{\mathrm{W}}\,\overline{\Psi}^{(2)}_{\mathrm{W}}\,,\\ H^- &= -\tfrac{1}{2}\partial\overline{\varphi}_{\mathrm{W}} - \tfrac{1}{2}(k+1)\partial\varphi_{\mathrm{W}} - \tfrac{1}{2}\Psi^{(1)}_{\mathrm{W}}\,\overline{\Psi}^{(1)}_{\mathrm{W}} - \tfrac{1}{2}\Psi^{(2)}_{\mathrm{W}}\,\overline{\Psi}^{(2)}_{\mathrm{W}} + \gamma_{\mathrm{W}}\,\beta_{\mathrm{W}}\,,\\ F_1 &= -\overline{\Psi}^{(2)}_{\mathrm{W}}\,\gamma_{\mathrm{W}} + \partial\varphi_{\mathrm{W}}\,\overline{\Psi}^{(1)}_{\mathrm{W}} - \partial\overline{\Psi}^{(1)}_{\mathrm{W}}\,,\\ F_2 &= \Psi^{(2)}_{\mathrm{W}}\,\gamma_{\mathrm{W}}\,\beta_{\mathrm{W}} - \partial\overline{\varphi}_{\mathrm{W}}\,\Psi^{(2)}_{\mathrm{W}} - (k+1)\Psi^{(1)}_{\mathrm{W}}\,\gamma_{\mathrm{W}} - \Psi^{(2)}_{\mathrm{W}}\,\Psi^{(1)}_{\mathrm{W}}\,\overline{\Psi}^{(1)}_{\mathrm{W}} + k\partial\Psi^{(2)}_{\mathrm{W}}\,,\\ F_{12} &= -\gamma_{\mathrm{W}}\,\gamma_{\mathrm{W}}\,\beta_{\mathrm{W}} + \partial\overline{\varphi}_{\mathrm{W}}\,\gamma_{\mathrm{W}} + (k+1)\partial\varphi_{\mathrm{W}}\,\gamma_{\mathrm{W}} - k\partial\gamma_{\mathrm{W}} + \Psi^{(1)}_{\mathrm{W}}\,\overline{\Psi}^{(1)}_{\mathrm{W}}\,\gamma_{\mathrm{W}} + \Psi^{(2)}_{\mathrm{W}}\,\overline{\Psi}^{(2)}_{\mathrm{W}}\,\gamma_{\mathrm{W}}\\ &\quad + \Psi^{(2)}_{\mathrm{W}}\,\partial\overline{\Psi}^{(1)}_{\mathrm{W}} - \partial\varphi_{\mathrm{W}}\,\Psi^{(2)}_{\mathrm{W}}\,\overline{\Psi}^{(1)}_{\mathrm{W}}\,.\end{aligned} \tag{6.9}$$

After the substitution (6.7), the currents (6.9) become the respective $s\ell(2|1)$ currents (6.3) with the $N=2$ matter generators taken in the bosonized form (6.6).

Furthermore, the standard Wakimoto screening operators [25, 18, 16] map under (6.7) into the $N=2$ *matter* screenings [38, 39]:

$$\begin{aligned}\Psi^{(1)}_{\mathrm{W}}\,e^{\varphi_{\mathrm{W}}} &= \overline{\chi}\,e^{X},\\ \Big(\overline{\Psi}^{(1)}_{\mathrm{W}}\,\beta_{\mathrm{W}} - (k+1)\overline{\Psi}^{(2)}_{\mathrm{W}}\Big)\,e^{\frac{1}{k+1}\overline{\varphi}_{\mathrm{W}}} &= \chi\,e^{\frac{1}{k+1}\overline{X}}.\end{aligned} \tag{6.10}$$

Similarly, the bosonic screening operator [25]

$$\Big(\partial\overline{\varphi}_{\mathrm{W}}\,\beta_{\mathrm{W}} + 3(k+1)^2\,\overline{\Psi}^{(2)}_{\mathrm{W}}\,\Psi^{(1)}_{\mathrm{W}} + (k+1)\Big[3\Psi^{(1)}_{\mathrm{W}}\,\overline{\Psi}^{(1)}_{\mathrm{W}}\,\beta_{\mathrm{W}} - 2\partial\varphi_{\mathrm{W}}\,\beta_{\mathrm{W}} + \partial\beta_{\mathrm{W}}\Big]\Big)\,\beta_{\mathrm{W}}^{-2-k}\,e^{-\overline{\varphi}_{\mathrm{W}}-(k+1)\varphi_{\mathrm{W}}} \tag{6.11}$$

evaluates under (6.7) as

$$\Big(\partial\overline{X} - 3(k+1)\chi\,\overline{\chi} - 2(k+1)\partial X\Big)\,e^{-\overline{X}-(k+1)X}, \tag{6.12}$$

thus reproducing the third screening in the bosonized matter theory (6.6). That the Wakimoto screening operators 'localize' in the matter sector under the mapping to the new representation appears to be a general fact (cf. [23]), and might be used as an alternative *definition* of the matter sector.

It should be pointed out that the screening charges written in terms of χ, $\overline{\chi}$, $\partial\overline{X}$ and ∂X imply a specific free-field realization (6.6) of the $N=2$ matter sector, which we only assume for the purposes of establishing a relation to the Wakimoto representation. Using a particular free-field realization is a matter of choice, and in that sense is not 'intrinsic', as is stressed in the $N=2$ case by the existence of a different free-field realization of the algebra (A.3), which we are now going to consider.

Now, consider an alternative realization of the $N=2$ matter [40, 16] in terms of a fermionic bc-system $\overline{\omega}\,\omega$, a 'Liouville' current $\partial\mathcal{F}$ and a free boson with the energy-momentum tensor $T_0 = \frac{1}{2}\partial\mathcal{X}\,\partial\mathcal{X} - (\frac{1}{2}\mathsf{a}_- +$

$\mathsf{a}_+)\partial\partial\mathcal{X}$ (with central charge $d_0 = 13 - 6(k+1) - \frac{6}{k+1}$, hence $k \neq -1$). This realization reads

$$\begin{aligned}
T_{\rm m} &= T_0 - \tfrac{1}{2}\partial\mathcal{F}\,\partial\mathcal{F} + \mathsf{a}_+\partial\partial\mathcal{F} - \tfrac{3}{2}\,\omega\,\partial\overline{\omega} - \tfrac{1}{2}\,\partial\omega\,\overline{\omega}\,,\\
H_{\rm m} &= \mathsf{a}_-\partial\mathcal{F} + \omega\,\overline{\omega}\,,\\
\overline{G}_{\rm m} &= \overline{\omega}\,T_0 - \tfrac{1}{2}\,\overline{\omega}\,\partial\mathcal{F}\,\partial\mathcal{F} + (\mathsf{a}_+ + \tfrac{1}{2}\mathsf{a}_-)\,\overline{\omega}\,\partial\partial\mathcal{F} + \tfrac{1}{2}(1-\mathsf{a}_-^2)\partial^2\,\overline{\omega} - \partial\,\overline{\omega}\,\overline{\omega}\,\omega + \mathsf{a}_-\partial\,\overline{\omega}\,\partial\mathcal{F}\,,\\
G_{\rm m} &= \omega\,,
\end{aligned} \tag{6.13}$$

$$\mathsf{a}_+ = \frac{1}{\sqrt{2(k+1)}}\,, \qquad \mathsf{a}_- = -\frac{1}{\mathsf{a}_+}\,.$$

The Wakimoto fields with the same operator products as in (6.8) can now be constructed in the following way:

$$\begin{aligned}
\beta_{\rm W} &= e^{\frac{2}{k}(U-F)}\,,\\
\gamma_{\rm W} &= \Big(\partial F - \sqrt{\tfrac{k+1}{2}}\,\partial\mathcal{X} + \tfrac{1}{2}\psi\,\overline{\psi} + \sqrt{\tfrac{k+1}{2}}\,\overline{\psi}\,\partial\mathcal{F}\,\overline{\omega} - \sqrt{\tfrac{k+1}{2}}\,\overline{\psi}\,\partial\mathcal{X}\,\overline{\omega} + (k+1)\overline{\psi}\,\partial\overline{\omega} + \tfrac{1}{2}\omega\,\overline{\omega}\Big)\,e^{-\frac{2}{k}(U-F)}\,,\\
\partial\varphi_{\rm W} &= \tfrac{1}{k}\,\partial F - \tfrac{1}{k}\,\partial U + \frac{1}{\sqrt{2(k+1)}}\,\partial\mathcal{F} - \frac{1}{\sqrt{2(k+1)}}\,\partial\mathcal{X}\,,\\
\partial\overline{\varphi}_{\rm W} &= -\tfrac{k+1}{k}\partial U + \tfrac{k+1}{k}\partial F - \sqrt{\tfrac{k+1}{2}}\,\partial\mathcal{F} - \sqrt{\tfrac{k+1}{2}}\,\partial\mathcal{X} + (k+1)\overline{\psi}\,\partial\overline{\omega} + (k+1)\partial\overline{\psi}\,\overline{\omega}\,,\\
\overline{\Psi}^{(1)}_{\rm W} &= \Big(\omega + \sqrt{\tfrac{k+1}{2}}\,\overline{\psi}\,\partial\mathcal{F} - \sqrt{\tfrac{k+1}{2}}\,\overline{\psi}\,\partial\mathcal{X} - (k+1)\,\partial\overline{\psi}\Big)\,e^{-\frac{1}{k}(U-F)}\,,\\
\Psi^{(1)}_{\rm W} &= -\overline{\omega}\,e^{\frac{1}{k}(U-F)}\,,\\
\overline{\Psi}^{(2)}_{\rm W} &= \overline{\psi}\,e^{\frac{1}{k}(U-F)}\,, \qquad\qquad \Psi^{(2)}_{\rm W} = \Big(\psi - \sqrt{\tfrac{k+1}{2}}\,\partial\mathcal{F}\,\overline{\omega} + \sqrt{\tfrac{k+1}{2}}\,\partial\mathcal{X}\,\overline{\omega} - (k+1)\partial\overline{\omega}\Big)\,e^{-\frac{1}{k}(U-F)}\,.
\end{aligned} \tag{6.14}$$

As before, these formulae provide a bridge between our realization (6.3) and the Wakimoto representation of $s\ell(2|1)$; however, the representation (6.3) (with the $N\!=\!2$ matter generators expressed as in (6.13)) will be reproduced if we start with the '*second*' Wakimoto representation of ref. [34], which can be written as[9]

$$\begin{aligned}
E_1 &= \Psi^{(2)}_{\rm W}\,\beta_{\rm W} - (k+1)\,\partial\varphi_{\rm W}\,\Psi^{(1)}_{\rm W} - (k+1)\,\partial\Psi^{(1)}_{\rm W}\,, \qquad E_2 = \overline{\Psi}^{(2)}_{\rm W}\,, \qquad E_{12} = \beta_{\rm W}\,,\\
H^+ &= -\tfrac{1}{2}\partial\overline{\varphi}_{\rm W} + \tfrac{1}{2}(k+1)\partial\varphi_{\rm W} - \tfrac{1}{2}\Psi^{(1)}_{\rm W}\,\overline{\Psi}^{(1)}_{\rm W} + \tfrac{1}{2}\Psi^{(2)}_{\rm W}\,\overline{\Psi}^{(2)}_{\rm W}\,,\\
H^- &= -\tfrac{1}{2}\partial\overline{\varphi}_{\rm W} - \tfrac{1}{2}(k+1)\partial\varphi_{\rm W} - \tfrac{1}{2}\Psi^{(1)}_{\rm W}\,\overline{\Psi}^{(1)}_{\rm W} - \tfrac{1}{2}\Psi^{(2)}_{\rm W}\,\overline{\Psi}^{(2)}_{\rm W} + \gamma_{\rm W}\,\beta_{\rm W}\,,\\
F_1 &= -\overline{\Psi}^{(2)}_{\rm W}\,\gamma_{\rm W} + \overline{\Psi}^{(1)}_{\rm W}\,,\\
F_2 &= \Psi^{(2)}_{\rm W}\,\gamma_{\rm W}\,\beta_{\rm W} - \partial\overline{\varphi}_{\rm W}\,\Psi^{(2)}_{\rm W} - (k+1)\,\partial\varphi_{\rm W}\,\Psi^{(1)}_{\rm W}\,\gamma_{\rm W} - \Psi^{(2)}_{\rm W}\,\Psi^{(1)}_{\rm W}\,\overline{\Psi}^{(1)}_{\rm W} - (k+1)\,\partial\Psi^{(1)}_{\rm W}\,\gamma_{\rm W} + k\partial\Psi^{(2)}_{\rm W}\,,\\
F_{12} &= -\gamma_{\rm W}\gamma_{\rm W}\beta_{\rm W} + \partial\overline{\varphi}_{\rm W}\gamma_{\rm W} + (k+1)\partial\varphi_{\rm W}\gamma_{\rm W} - k\partial\gamma_{\rm W} + \Psi^{(1)}_{\rm W}\overline{\Psi}^{(1)}_{\rm W}\gamma_{\rm W} - \Psi^{(2)}_{\rm W}\overline{\Psi}^{(1)}_{\rm W} + \Psi^{(2)}_{\rm W}\overline{\Psi}^{(2)}_{\rm W}\gamma_{\rm W}
\end{aligned} \tag{6.15}$$

It is amusing to see how the various pieces of the $N\!=\!2$ matter generators (6.13) find each other when eqs. (6.14) are inserted into (6.15).

Thus, allowing the $N = 2$ matter generators to be bosonized through free fields, we can relate the corresponding Fock space image of the representation (6.3) to the respective Wakimoto representation. Not surprisingly [16], the two different bosonizations of the $N\!=\!2$ matter are thus related to the respective inequivalent simple root systems of $s\ell(2|1)$. By restricting to $osp(1|2)$ and $s\ell(2)$, one can work out similar relations for representations of these algebras.

---

[9] Certain rescalings of the type of eq. (2.6) would slightly simplify the expressions for the Wakimoto fields, at the same time introducing factors in front of some terms in (6.15); in the present normalization the formulae (6.15) have as much as possible (e.g., the Cartan subalgebra) in common with the 'first' Wakimoto representation (6.9).

7 Concluding remarks

We have shown that any non-critical $N=2$ string in the conformal gauge reformulates as the $s\ell(2|1)$ theory tensored with two bosonic and two fermionic 'Drinfeld–Sokolov' ghost systems. The construction of the $s\ell(2|1)$ currents appears to be 'rigid', admitting no free parameters, modulo the above-mentioned automorphisms of the defining operator products (it is a useful exercise to lift the spectral flow transform [31, 32] to the $s\ell(2|1)$ algebra).

The present construction may provide new group-theoretical '*explanations*' for a number of hidden supersymmetry algebras that arise in non-critical string theories.

An important problem left for the future investigation is that of a precise comparison of the *physical states* (i.e., the BRST cohomologies) of the $s\ell(2|1)$ theory and the $N=2$ string. The problem may be feasible by the methods developed in conformal field theory. In the $N=2$ matter sector, we have just the corresponding *Verma* modules, hence no need for screening charges to project from the Fock modules. However, the two $\beta\gamma$ systems that participate in the construction (2.7)–(2.12) are potentially more interesting from the cohomological point of view: as we have noted, the $s\ell(2|1)$ currents depend only on the respective $\beta\gamma$ *currents*, which excludes, by the well-known argument (see [41] and numerous subsequent applications of the idea), the zero mode of a spin-0 fermion that participates in the bosonization of the $\beta\gamma$ system.

One may wonder how special the realization (2.7)–(2.12) of the $s\ell(2|1)$ algebra is. A related question is whether one can reconstruct the non-critical $N=2$-string fields from an 'abstract' $s\ell(2|1)$ algebra *and* the Drinfeld–Sokolov ghosts (we have seen in (2.14) that only with these ghosts added do the energy-momentum tensors match and thus there is a chance to recover the $N=2$ string). The $N=2$ matter sector can indeed be recovered, in quite abstract terms, via the hamiltonian reduction. The two bosonic and two fermionic Drinfeld–Sokolov ghost systems can roughly (in fact, modulo some mixing with the other free fields) be traded for the same number of bosonic and fermionic $N=2$ *string* ghosts. The problem is therefore concentrated in the $N=2$ Liouville sector (2.4). The current $\overline{\psi}\psi$ could then be found from the second equation in (2.8) (assuming the matter $U(1)$ current $H_{\rm m}$ known from the hamiltonian reduction). This leaves us with the first equation in (2.8), which does not, however, allow us to determine $\partial\overline{\phi}$ and $\partial\phi$ in terms of the given $s\ell(2|1)$ currents and the other ingredients which we assumed known: the matter $U(1)$ current $H_{\rm m}$ and the ghost currents $B\,C$, $\mathsf{B}\,\Gamma$, $\widetilde{\mathsf{B}}\,\widetilde{\Gamma}$, and $\mathcal{B}\,\mathcal{C}$; only the combination $-\partial\overline{\phi}+(k+3)\partial\phi$ can be determined. That finding $\partial\overline{\phi}$ and $\partial\phi$ and hence all the $N=2$ string ingredients should require some special assumptions is also supported indirectly by the fact that the $N=2$ string has an $N=4$ symmetry [17]; if we had been able to identify the $N=2$ non-critical string in a general representation of $s\ell(2|1)\otimes(\text{DS ghosts})$, we would have had to conclude that the $N=4$ generators could be constructed in any such representation as well. Anyway, it would be interesting to find the mildest requirements that would make that possible.

As we have seen, the module in which the $s\ell(2|1)$ currents (2.7)–(2.12) are represented can be constructed by tensoring the $N=2$ matter *Verma* module with free field Fock spaces. This allows us to expect that the $s\ell(2|1)$ singular vectors would evaluate as the $N=2$ superconformal singular vectors (thus it happens in the $s\ell(2)$ case [40, 23, 33]). However, the effects due to the different types of the $N=2$ [21] and $s\ell(2|1)$ singular vectors and the *sub*singular vectors of these algebras will have to be accounted for, which appears to be an interesting problem.

Acknowledgements. I am grateful to P. Bowcock, A. Taormina and A. Tseytlin for stimulating my interest in $N=2$ strings and for very interesting discussions. Extensive computations of the operator product expansions have been done using the MATHEMATICA package `OPEdefs.m` by K. Thielemans [42]. This work was supported in part by the RFFI grant 96-02-16117 and by the European Community grant INTAS-93-2058.

Appendix

The $s\ell(2|1)$ algebra consists of four bosonic currents E_{12}, F_{12}, H^- and H^+, and four fermionic ones E_1, E_2, F_1 and F_2. The non-vanishing OPEs read,

$$
\begin{array}{rclrcl}
H^{\pm}(z)\,H^{\pm}(w) & = & \mp\dfrac{k/2}{(z-w)^2}\,, & E_{12}(z)\,F_{12}(w) & = & \dfrac{k}{(z-w)^2}+2\dfrac{H^-}{z-w}\,,\\
H^-(z)\,E_{12}(w) & = & \dfrac{E_{12}}{z-w}\,, & H^-(z)\,F_{12}(w) & = & \dfrac{-F_{12}}{z-w}\,,\\
E_{12}(z)\,F_1(w) & = & \dfrac{E_2}{z-w}\,, & E_{12}(z)\,F_2(w) & = & -\dfrac{E_1}{z-w}\,,\\
F_{12}(z)\,E_1(w) & = & -\dfrac{F_2}{z-w}\,, & F_{12}(z)\,E_2(w) & = & \dfrac{F_1}{z-w}\,,\\
H^{\pm}(z)\,E_1(w) & = & \frac{1}{2}\dfrac{E_1}{z-w}\,, & H^{\pm}(z)\,F_1(w) & = & -\frac{1}{2}\dfrac{F_1}{z-w}\,,\\
H^{\pm}(z)\,E_2(w) & = & \mp\frac{1}{2}\dfrac{E_2}{z-w}\,, & H^{\pm}(z)\,F_2(w) & = & \pm\frac{1}{2}\dfrac{F_2}{z-w}\,,\\
E_1(z)\,F_1(w) & = & -\dfrac{k}{(z-w)^2}+\dfrac{H^+-H^-}{z-w}\,, & E_2(z)\,F_2(w) & = & \dfrac{k}{(z-w)^2}+\dfrac{H^++H^-}{z-w}\,,\\
E_1(z)\,E_2(w) & = & \dfrac{E_{12}}{z-w}\,, & F_1(z)\,F_2(w) & = & \dfrac{F_{12}}{z-w}
\end{array}
\tag{A.1}
$$

all the other OPEs being regular. The Sugawara energy-momentum tensor reads

$$
T_{\rm Sug}=\tfrac{1}{k+1}\Bigl(H^-\,H^- - H^+\,H^+ + \tfrac{1}{2}E_{12}\,F_{12}+\tfrac{1}{2}F_{12}\,E_{12}+\tfrac{1}{2}E_1\,F_1-\tfrac{1}{2}F_1\,E_1-\tfrac{1}{2}E_2\,F_2+\tfrac{1}{2}F_2\,E_2\Bigr)
\tag{A.2}
$$

The $N=2$ matter, which is one of the ingredients to construct the $s\ell(2|1)$ currents, is defined by

$$
\begin{array}{rclrcl}
T_{\rm m}(z)\,T_{\rm m}(w) & = & \dfrac{c_{\rm m}/2}{(z-w)^4}+2\dfrac{T_{\rm m}(w)}{(z-w)^2}+\dfrac{\partial T_{\rm m}(w)}{z-w}\,, & & & \\
H_{\rm m}(z)\,H_{\rm m}(w) & = & \dfrac{c_{\rm m}/3}{(z-w)^2}\,, & T_{\rm m}(z)\,H_{\rm m}(w) & = & \dfrac{H_{\rm m}(w)}{(z-w)^2}+\dfrac{\partial H_{\rm m}(w)}{z-w}\,,\\
T_{\rm m}(z)\,G_{\rm m}(w) & = & \frac{3}{2}\dfrac{G_{\rm m}(w)}{(z-w)^2}+\dfrac{\partial G_{\rm m}(w)}{z-w}\,, & T_{\rm m}(z)\,\overline{G}_{\rm m}(w) & = & \frac{3}{2}\dfrac{\overline{G}_{\rm m}(w)}{(z-w)^2}+\dfrac{\partial\overline{G}_{\rm m}(w)}{z-w}\,,\\
H_{\rm m}(z)\,G_{\rm m}(w) & = & \dfrac{G_{\rm m}(w)}{z-w}\,, & H_{\rm m}(z)\,\overline{G}_{\rm m}(w) & = & -\dfrac{\overline{G}_{\rm m}(w)}{z-w}\,,\\
\overline{G}_{\rm m}(z)\,G_{\rm m}(w) & = & \dfrac{c_{\rm m}/3}{(z-w)^3}-\dfrac{H_{\rm m}(w)}{(z-w)^2}+\dfrac{T_{\rm m}(w)-\frac{1}{2}\partial H_{\rm m}(w)}{z-w} & & &
\end{array}
\tag{A.3}
$$

For completeness, let us also list the non-vanishing $osp(1|2)$ OPEs:

$$\begin{array}{rclcrcl}
I^{+}(z)I^{-}(w) & = & \dfrac{k}{(z-w)^2}+\dfrac{2I^0}{z-w}, & & I^0(z)I^0(w) & = & \dfrac{k/2}{(z-w)^2},\\
I^{+\frac{1}{2}}(z)I^{-\frac{1}{2}}(w) & = & \dfrac{-2k}{(z-w)^2}-2\dfrac{I^0}{z-w}, & & I^0(z)I^{\pm}(w) & = & \pm\dfrac{I^{\pm}}{(z-w)}\\
I^0(z)I^{\pm\frac{1}{2}}(w) & = & \pm\frac{1}{2}\dfrac{I^{\pm\frac{1}{2}}}{z-w}, & & I^{\pm}(z)I^{\mp\frac{1}{2}}(w) & = & -\dfrac{I^{\pm\frac{1}{2}}}{z-w},\\
I^{\pm\frac{1}{2}}(z)I^{\pm\frac{1}{2}}(w) & = & \mp2\dfrac{I^{\pm}}{z-w} & & & &
\end{array} \tag{A.4}$$

References

[1] M. Ademollo, L. Brink, A. D'Adda, R. D'Auria, E. Napolitano, S. Sciuto, E. Del Guidice, P. Di Vecchia, S. Ferrara, F. Gliozzi, R. Musto, and R. Pettorino, Phys. Lett. B62 (1976) 105;
M. Ademollo, L. Brink, A. D'Adda, R. D'Auria, E. Napolitano, S. Sciuto, E. Del Guidice, P. Di Vecchia, S. Ferrara, F. Gliozzi, R. Musto, R. Pettorino, and J.H. Schwarz, Nucl. Phys. B111 (1976) 77.

[2] N. Marcus, *A tour through $N = 2$ strings*, talk at the Rome String Theory Workshop, 1992, hep-th/9211059.

[3] E.S. Fradkin and A.A. Tseytlin, Phys. Lett. B106 (1981) 63; Phys. Lett. B162 (1985) 295.

[4] A. D'Adda and F. Lizzi, Phys. Lett. B191 (1987) 85.

[5] S. Mathur and S. Mukhi, Phys. Rev. D36 (1987) 465; Nucl. Phys. B302 (1988) 130.

[6] H. Ooguri and C. Vafa Nucl. Phys. B361 (1991) 469; Nucl. Phys. B367 (1991) 83.

[7] O. Lechtenfeld, *Integration Measure and Spectral Flow in the Critical $N=2$ String*, hep-th/9512189.

[8] D. Kutasov and E. Martinec, *New Principles for String/Membrane Unification*, hep-th/9602049.

[9] L. Brink and J. Schwarz, Nucl. Phys. B121 (1977) 285.

[10] N. Berkovits and C. Vafa, Mod. Phys. Lett. A9 (1994) 653.

[11] J.M. Figueroa-O'Farrill, Phys. Lett. B321 (1994) 344; Nucl. Phys. B432 (1994) 404.

[12] N. Ohta and J.L. Petersen, Phys. Lett. B325 (1994) 67.

[13] F. Bastianelli, N. Ohta and J.L. Petersen, Phys. Lett. B327 (1994) 35.

[14] N. Berkovits and N. Ohta, Phys. Lett. B334 (1994) 72.

[15] B. Gato-Rivera and A.M. Semikhatov, Phys. Lett. B293 (1992) 72, Theor. Math. Phys. 95 (1993) 536.

[16] M. Bershadsky, W. Lerche, D. Nemeschansky, and N.P. Warner, Nucl. Phys. B401 (1993) 304.

[17] A. Boresch, K. Landsteiner, W. Lerche and A. Sevrin, Nucl. Phys. B436 (1995) 609.

[18] K. Ito and H. Kanno, Mod. Phys. Lett. A9 (1994) 1377.

[19] E. Ragoucy, A. Sevrin and P. Sorba, *Strings from $N = 2$ Gauged Wess-Zumino-Witten Models*, hep-th/9511049.

[20] K. Landsteiner, W. Lerche and A. Sevrin, Phys. Lett. B352 (1995) 286.

[21] A.M. Semikhatov and I.Yu. Tipunin, *All Singular Vectors of the $N=2$ Superconformal Algebra via the Algebraic Continuation Approach*, hep-th/9604176.

[22] B.L. Feigin, A.M. Semikhatov, and I.Yu. Tipunin, *Equivalence between Chain Categories of Representations of Affine $s\ell(2)$ and $N = 2$ Superconformal Algebras*, hep-th/9701043.

[23] A.M. Semikhatov, Mod. Phys. Lett. A9 (1994) 1867.

[24] C.M. Hull and A.M. Semikhatov, unpublished.

[25] M. Bershadsky and H. Ooguri, Phys. Lett. B229 (1989) 374.

[26] E. Verlinde and H. Verlinde, Nucl. Phys. B352 (1991) 1234;
R. Dijkgraaf, E. Verlinde, and H. Verlinde, Nucl. Phys. B352 (1991) 59.

[27] F. David, Mod. Phys. Lett. A3 (1988) 1651.

[28] J. Distler and H. Kawai, Nucl. Phys. B321 (1989) 509.

[29] A.M. Polyakov, Mod. Phys. Lett. A2 (1987) 893.

[30] V.G. Knizhnik, A.M. Polyakov, and A.B. Zamolodchikov, Mod. Phys. Lett. A3 (1988) 819.

[31] A. Schwimmer and N. Seiberg, Phys. Lett. B184 (1987) 191.

[32] W. Lerche, C. Vafa, and N.P. Warner, Nucl. Phys. B324 (1989) 427.

[33] B. Gato-Rivera and J.-I. Rosado, Phys. Lett. B346 (1995) 63.

[34] P. Bowcock, R.L. Koktava, and A. Taormina, *Free field Representations for the Affine Superalgebra* $sl\widehat{(2|1)}$ *and noncritical* $N=2$ *strings*, hep-th/9606015.

[35] G. Mussardo, G. Sotkov, and M. Stanishkov, Int. J. Mod. Phys. A4 (1989) 1135.

[36] N. Ohta and H. Suzuki, Nucl. Phys. B332 91990) 146.

[37] K. Ito, Nucl. Phys. B370 (1992) 123.

[38] P. Di Vecchia, J.L. Petersen, and H.B. Zheng, Phys. Lett. B162 (1985) 327.

[39] M. Yu and H.B. Zheng, Nucl. Phys. B288 (1987) 275.

[40] B. Gato-Rivera and A.M. Semikhatov, Nucl. Phys. B408 (1993) 133.

[41] D.H. Friedan, E.J. Martinec, and S.H. Shenker, Nucl. Phys. B271 (1986) 93.

[42] K. Thielemans, Int. J. Mod. Phys. C2, No. 3, (1991) 787.